arXiv:1204.2748v1 [math.AP] 12 Apr 2012

HOMOGENIZATION OF WEAKLY COUPLED SYSTEMS OF HAMILTON–JACOBI EQUATIONS WITH FAST SWITCHING RATES

HIROYOSHI MITAKE AND HUNG V. TRAN

Dedicated to Professor H. Ishii on the occasion of his 65th birthday

Abstract. We consider homogenization for weakly coupled systems of Hamilton–Jacobi equations with fast switching rates. The fast switching rate terms force the solutions converge to the same limit, which is a solution of the effective equation. We discover the appearance of the initial layers, which appear naturally when we consider the systems with different initial data and analyze them rigorously. In particular, we obtain matched asymptotic solutions of the systems and rate of convergence. We also investigate properties of the effective Hamiltonian of weakly coupled systems and show some examples which do not appear in the context of single equations.

Contents

1. Introduction 2
1.1. Background: Randomly Switching Cost Problems 2
1.2. Main Results 3
2. Homogenization Results 7
3. Initial layers and Rate of convergence 9
3.1. Inner solutions, Outer solutions, and Matched solutions 9
3.2. Rate of convergence to matched solutions 11
4. Properties of effective Hamiltonians 14
4.1. Elementary properties 14
4.2. Representation formulas for the effective Hamiltonian 16
4.3. Flat parts and Non-flat parts near the origin 19
5. Generalization 26
6. Dirichlet Problems 27

Date: March 2, 2013.

2010 *Mathematics Subject Classification.* 35B27, 35F55, 49L25.

Key words and phrases. Homogenization; Hamilton–Jacobi Equations; Weakly Coupled Systems; Effective Hamiltonians; Switching Cost Problems; Piecewise-deterministic Markov processes; Viscosity Solutions.

This work was partially done while the first author visited Mathematics Department, University of California, Berkeley.

7. Appendix 30
References 34

1. Introduction

In this paper we study the behavior, as $\varepsilon(>0)$ tends to 0, of the viscosity solutions $(u_1^\varepsilon, u_2^\varepsilon)$ of the following weakly coupled systems of Hamilton–Jacobi equations

$$(\mathrm{C}_\varepsilon)\qquad \begin{cases} (u_1^\varepsilon)_t + H_1(\dfrac{x}{\varepsilon}, Du_1^\varepsilon) + \dfrac{c_1}{\varepsilon}(u_1^\varepsilon - u_2^\varepsilon) = 0 & \text{in } \mathbb{R}^n \times (0,T), \\ (u_2^\varepsilon)_t + H_2(\dfrac{x}{\varepsilon}, Du_2^\varepsilon) + \dfrac{c_2}{\varepsilon}(u_2^\varepsilon - u_1^\varepsilon) = 0 & \text{in } \mathbb{R}^n \times (0,T), \\ u_i^\varepsilon(x,0) = f_i(x) & \text{on } \mathbb{R}^n \text{ for } i = 1,2, \end{cases}$$

where $T > 0$, c_1, c_2 are given positive constants and the Hamiltonians $H_i(\xi, p) : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ are given continuous functions for $i = 1, 2$, which are assumed throughout the paper to satisfy the followings.

(A1) The functions H_i are uniformly coercive in the ξ-variable, i.e.,

$$\lim_{r\to\infty} \inf\{H_i(\xi, p) \mid \xi \in \mathbb{R}^n, |p| \geq r\} = \infty.$$

(A2) The functions $\xi \mapsto H_i(\xi, p)$ are $\mathbb{T}^n$-periodic, i.e., $H_i(\xi + z, p) = H_i(\xi, p)$ for any $\xi, p \in \mathbb{R}^n$, $z \in \mathbb{Z}^n$ and $i = 1, 2$.

The functions f_i are given continuously differentiable functions on $\mathbb{R}^n$ with $\|Df_i\|_{L^\infty(\mathbb{R}^n)}$ are bounded for $i = 1, 2$, respectively. Here u_i^ε are the real-valued unknown functions on $\mathbb{R}^n \times [0, T]$ and $(u_i^\varepsilon)_t := \partial u_i^\varepsilon/\partial t, Du_i^\varepsilon := (\partial u_i^\varepsilon/\partial x_1, \dots, \partial u_i^\varepsilon/\partial x_n)$ for $i = 1, 2$, respectively. We are dealing only with viscosity solutions of Hamilton–Jacobi equations in this paper and thus the term "viscosity" may be omitted henceforth.

1.1. **Background: Randomly Switching Cost Problems.** System (C_ε) arises as the dynamic programming for the optimal control of the system whose states are governed by certain ODEs, subject to random changes in the dynamics: the system randomly switches at a fast rate $1/\varepsilon$ among the two states. See [13, 14, 16] for instance. Also see [26, 7] for another switching cost problems. In order to explain the background more precisely, we assume in addition that the Hamiltonians H_i are convex in p here. We define the functions $u_i^\varepsilon : \mathbb{R}^n \times [0, T] \to \mathbb{R}$ by

$$u_i^\varepsilon(x,t) := \inf\Big\{\mathbb{E}_i\Big(\int_0^t L_{\nu^\varepsilon(s)}(\frac{\eta(s)}{\varepsilon}, -\dot\eta(s))\, ds + f_{\nu^\varepsilon(t)}(\eta(t))\Big)\Big\}, \tag{1.1}$$

where $L_i : \mathbb{R}^{2n} \to \mathbb{R} \cup \{+\infty\}$ are the Fenchel-Legendre transform of H_i, i.e., $L_i(\xi, q) := \sup_{p\in\mathbb{R}^n}(p \cdot q - H_i(\xi, p))$ for all $(\xi, q) \in \mathbb{R}^{2n}$ and the infimum is taken over $\eta \in \mathrm{AC}([0, t], \mathbb{R}^n)$ such that $\eta(0) = x$. Here $\mathrm{AC}([0, t], \mathbb{R}^n)$ denotes the set of absolutely continuous functions

with value in $\mathbb{R}^n$ and $\mathbb{E}_i$ denotes the expectation of a process with $\nu^\varepsilon(0) = i$ where ν is a $\{1,2\}$-valued process which is a continuous-time Markov chain such that

$$\mathbb{P}\big(\nu^\varepsilon(s+\Delta s) = j \mid \nu^\varepsilon(s) = i\big) = \frac{c_i}{\varepsilon}\Delta s + o(\Delta s) \text{ as } \Delta s \to 0 \text{ for } i \neq j, \tag{1.2}$$

where $o : [0,\infty) \to [0,\infty)$ is a function which satisfies $o(r)/r \to 0$ as $r \to 0$. Formula (1.1) is basically the optimal control formula for the solution of (C_ε), where the random switchings among the two states are governed by (1.2).

We first give a formal proof that $(u_1^\varepsilon, u_2^\varepsilon)$ given by (1.1) is a solution of (C_ε). The rigorous derivation will be proved in Appendix by using the dynamic programming principle. We suppose that $u_i \in C^1(\mathbb{R}^n \times [0,T])$ here. Set $u(x,i,t) := u_i(x,t)$ and $Y(s) := (\eta(s), \nu^\varepsilon(s))$ for $\eta \in \mathrm{AC}\,(\mathbb{R}^n)$ with $\eta(0) = x$ and let ν^ε be a Markov chain given by (1.2) with $\nu^\varepsilon(0) = i$. By Ito's formula for a jump process we have

$$\begin{aligned}
&\mathbb{E}_i\Big(u^\varepsilon(Y(t),0) - u^\varepsilon(Y(0),t)\Big)\\
=&\,\mathbb{E}_i\Big(\int_0^t -u_t^\varepsilon(Y(s),t-s) + Du^\varepsilon(Y(s),t-s)\cdot\dot\eta(s)\,ds\\
&\quad+\int_0^t \sum_{j=1}^2 \big(u^\varepsilon(\eta(s),j,s) - u^\varepsilon(\eta(s),\nu^\varepsilon(s),s)\big)\cdot\frac{c_{\nu^\varepsilon(s)}}{\varepsilon}\,ds\Big)\\
\geq&\,\mathbb{E}_i\Big(\int_0^t -u_t^\varepsilon(Y(s),t-s) - H_{\nu^\varepsilon(s)}(\frac{\eta}{\varepsilon}, Du^\varepsilon) - L_{\nu^\varepsilon(s)}(\frac{\eta}{\varepsilon}, -\dot\eta)\,ds\\
&\quad+\int_0^t \sum_{j=1}^2 \big(u^\varepsilon(\eta(s),j,s) - u^\varepsilon(\eta(s),\nu^\varepsilon(s),s)\big)\cdot\frac{c_{\nu^\varepsilon(s)}}{\varepsilon}\,ds\Big)\\
=&\,-\mathbb{E}_i\Big(\int_0^t L_{\nu^\varepsilon(s)}(\frac{\eta}{\varepsilon}, -\dot\eta)\,ds\Big).
\end{aligned}$$

Thus,

$$u^\varepsilon(x,i,t) \leq \mathbb{E}_i\Big(\int_0^t L_{\nu^\varepsilon(s)}(\frac{\eta}{\varepsilon}, -\dot\eta)\,ds + u^\varepsilon(Y(t),0)\Big).$$

In the above inequality, the equality holds if $-\dot\eta(s) \in D_p^- H_{\nu^\varepsilon(s)}(\eta(s)/\varepsilon, Du(Y(s),s))$, where $D_p^- H_i$ denotes the subdifferential of H_i with respect to the p-variable.

1.2. **Main Results.** There have been extensively many important results on the study of homogenization of Hamilton–Jacobi equations. The first general result is due to Lions, Papanicolaou, and Varadhan [27] who studied the cell problems together with the effective Hamiltonian and established homogenization results under quite general assumptions on Hamiltonians in the periodic setting. The next major contributions to the subject are due to Evans [16, 17] who introduced the perturbed test function method in the framework of viscosity solutions. The method then has been adapted to study so many different

homogenization problems that we cannot describe the complete list of references. We here only refer to the papers related to our work. Concordel [11, 12] achieved some first general results on the properties of the effective Hamiltonian concerning flat parts and non-flat parts. Afterwards Capuzzo-Dolceta and Ishii [9] combined the perturbed test functions with doubling variables methods to obtain the first results on the rate of convergence of u^ε to u. We refer to [5, 35] for some recent progress.

There have been some interesting results [33, 6, 8] on the study of homogenization for weakly coupled systems of Hamilton–Jacobi equations in the periodic settings or in the almost periodic settings. We refer the readers to [15, 23] for the complete theory of viscosity solutions for weakly coupled systems of Hamilton–Jacobi and Hamilton–Jacobi–Bellman equations. Since the maximum principle and comparison principle still hold, homogenization results can be obtained by using the perturbed test function method quite straightforwardly with some modifications. Let us call attention also to the new interesting direction on the large time behavior of weakly coupled sytems of Hamilton–Jacobi equations, which is related to homogenization through the cell problems. The authors [30], and Camilli, Ley, Loreti and Nguyen [7] obtained large time behavior results for some special cases but general cases still remain open.

Let us also refer to one of the main research directions in the study of homogenization, stochastic homogenization of Hamilton–Jacobi equations, which were first obtained by Souganidis [34], and Rezakhanlou and Tarver [31] independently. See [28, 25, 32, 29, 1] for more recent progress on the subject.

First we heuristically derive the behavior of solutions of (C_ε) as ε tends to 0. For simplicity, from now on, we *always* assume that $c_1 = c_2 = 1$. We consider the formal asymptotic expansions of solutions $(u_1^\varepsilon, u_2^\varepsilon)$ of (C_ε) of the form

$$u_i^\varepsilon(x,t) := u_i(x,t) + \varepsilon v_i(\frac{x}{\varepsilon}) + O(\varepsilon^2).$$

Set $\xi := x/\varepsilon$. Plugging this into (C_ε) and performing formal calculations, we achieve

$$(u_i)_t + \ldots + H_i(\xi, D_x u_i + D_\xi v_i + \cdots) + \frac{1}{\varepsilon}(u_i - u_j) + (v_i - v_j) + \cdots = 0,$$

where we take $i, j \in \{1,2\}$ such that $\{i,j\} = \{1,2\}$. The above expansion implies that $u_1 = u_2 =: u$. Furthermore, if we let $P = Du(x,t)$ then (v_1, v_2) is a $\mathbb{T}^n$-periodic solution of the following cell problem

$$(\mathrm{E}_P) \qquad \begin{cases} H_1(\xi, P + Dv_1(\xi,P)) + v_1(\xi,P) - v_2(\xi,P) = \overline{H}(P) & \text{in } \mathbb{R}^n, \\ H_2(\xi, P + Dv_2(\xi,P)) + v_2(\xi,P) - v_1(\xi,P) = \overline{H}(P) & \text{in } \mathbb{R}^n, \end{cases}$$

where $\overline{H}(P)$ is a unknown constant. Because of the $\mathbb{T}^n$-periodicity of the Hamiltonians H_i, we can also consider the above cell problem on the torus $\mathbb{T}^n$, which is equivalent to

consider it on $\mathbb{R}^n$ with $\mathbb{T}^n$-periodic solutions. By an argument similar to the classical one in [27], we have

Proposition 1.1 (Cell Problems). *For any $P \in \mathbb{R}^n$, there exists a unique constant $\overline{H}(P)$ such that* (E_P) *admits a $\mathbb{T}^n$-periodic solution $(v_1(\cdot, P), v_2(\cdot, P)) \in C(\mathbb{R}^n)^2$. We call $\overline{H}$ the effective Hamiltonian assosiated with (H_1, H_2).*

See also [4, 30, 7] for more details about the cell problems for weakly coupled systems.

Our main goal in this paper is threefold. First of all, we want to demonstrate that u_i^ε converge locally uniformly to the same limit u in $\mathbb{R}^n \times (0,T)$ for $i = 1, 2$ and u solves

$$u_t + \overline{H}(Du) = 0 \quad \text{in } \mathbb{R}^n \times (0,T).$$

This part is a rather standard part in the study of homogenization of Hamilton–Jacobi equations by using the perturbed test function method introduced by Evans [16] with some modifications. The only hard part comes from the fact that we do not have uniform bounds on the gradients of u_i^ε here because of the fast switching terms. We overcome this difficulty by introducing the barrier functions (see Lemma 2.1) and using the half-relaxed limits (see the proof of Theorem 1.2). The barrier functions furthermore give us the correct initial data for the limit u. Let $(u_1^\varepsilon, u_2^\varepsilon)$ be the solution of (C_ε) henceforth.

Theorem 1.2 (Homogenization Result). *The functions u_i^ε converge locally uniformly in $\mathbb{R}^n \times (0,T)$ to the same limit $u \in C^{0,1}(\mathbb{R}^n \times [0,T])$ as $\varepsilon \to 0$ for $i = 1, 2$ and u solves*

$$\begin{cases} u_t + \overline{H}(Du) = 0 & \text{in } \mathbb{R}^n \times (0,T) \\ u(x,0) = \overline{f}(x) := \dfrac{f_1(x) + f_2(x)}{2} & \text{on } \mathbb{R}^n. \end{cases} \tag{1.3}$$

Formula (1.1) of solutions of (C_ε) actually gives us an intuitive explanation about the effective initial datum $\overline{f}$. As we send ε to 0, the switching rate becomes very fast and processes have to jump randomly very quickly between the two states with equal probability as given by (1.2). (Note that we are assuming $c_1 = c_2 = 1$ now.) Therefore, it is relatively clear that $\overline{f}$ is the average of the given initial data f_i for $i = 1, 2$.

The second main part of this paper is the study of the initial layers appearing naturally in the problem as the initial data of u_i^ε and u are different in general. We first study the initial layers in a heuristic mode by finding inner and outer solutions, and using the matching asymptotic expansion method to identify matched solutions (see Section 3.1). We then combine the techniques of the matching asymptotic expansion method and of Capuzzo-Dolceta and Ishii [9] to obtain rigorously the rate of convergence result.

Theorem 1.3 (Rate of Convergence to Matched Solutions). *For each $T > 0$, there exists $C := C(T) > 0$ such that*

$$\|u_i^\varepsilon - m_i^\varepsilon\|_{L^\infty(\mathbb{R}^n \times [0,T])} \le C\varepsilon^{1/3} \text{ for } i = 1, 2,$$

where u is the solution of (1.3) *and*

$$m_i^\varepsilon(x,t) := u(x,t) + \frac{(f_i - f_j)(x)}{2} e^{-\frac{2t}{\varepsilon}} \tag{1.4}$$

with $j \in \{1,2\}$ such that $\{i,j\} = \{1,2\}$.

Finally, we study various properties of the effective Hamiltonian $\overline{H}$. It is always extremely hard to understand properties of the effective Hamiltonians even for single equations. Lions, Papanicolaou and Vadrahan [27] studied some preliminary properties of the effective Hamiltonians and pointed out a 1-dimensional example that $\overline{H}$ can be computed explicitly. After that, Concordel [11, 12] discovered some very interesting results related flat parts and non-flat parts of $\overline{H}$ for more general cases. Evans and Gomes [18] found some further properties on the strict convexity of $\overline{H}$ by using the weak KAM theory.

The properties of $\overline{H}$ for weakly coupled systems of Hamilton–Jacobi equations in this paper are even more complicated. In case $H_1 = H_2$, the effective Hamiltonian for the weakly coupled systems and the single equations are obviously same. Therefore, we can view the cases of single equations as special cases of the weakly coupled systems. However, in general, we cannot expect the effective Hamiltonians for weakly coupled systems to have similar properties like single equations' cases.

The first few results on flat parts and non-flat parts of $\overline{H}$ are generalizations to the ones discovered by Concordel [11, 12], and are proved by using different techniques, namely the min-max formulas which are derived in Section 4.2 and the constructions of appropriate subsolutions. On the other hand, we investigate other cases which show that the properties of the effective Hamiltonians for weakly coupled systems are widely different from those of the effective Hamiltonians for single equations. Theorems 4.14, 4.17, 4.18 , 4.20, which are ones of our main results, describe some rather new results which do not appear in the context of single Hamilton-Jacobi equations. Since the theorems are technical, we refer the readers to Section 4.3 for details.

We are grateful to L. C. Evans for his suggestion which leads us to this project. We thank G. Barles, D. Gomes, H. Ishii, T. Mikami, and F. Rezakhanlou for their fruitful discussions. We also thank S. Armstrong and P. E. Souganidis for letting us know about the coming result on stochastic homogenization of weakly coupled systems of Hamilton–Jacobi equations of B. Fehrman [19]. Fehrman [19] independently obtained interesting homogenization results of monotone systems of viscous Hamilton–Jacobi equations, which are similar to ours, with convex Hamiltonians in the stationary, ergodic setting by using the ideas of Armstrong and Souganidis [1, 2]. His work includes as well generalizations to other related systems.

The paper is organized as follows. In Section 2 we prove the homogenization result, Theorem 1.2. Section 3 devotes to the study of initial layers and rate of convergence. We derive inner solutions, outer solutions, and matched asymptotic solutions in a heuristic

mode and then prove Theorem 1.3. The properties of the effective Hamiltonian are studied in Section 4. We obtain its elementary properties in Section 4.1, the representation formulas in Section 4.2, and flat parts, non-flat parts near the origin in Section 4.3. In Section 5 we prove generalization results for systems of m equations for $m \geq 2$. We then prove also the homogenization result for Dirichlet problems and describe the differences of the effective data between Cauchy problems and Dirichlet problems in Section 6. Some lemmata concerning verifications of optimal control formulas for the Cauchy and Dirichlet problems are recorded in Appendix.

Notations. For $k \in \mathbb{N}$ and $A \subset \mathbb{R}^n$, we denote by $C(A)$, $C^{0,1}(A)$ and $C^k(A)$ the space of real-valued continuous, Lipschitz continuous and k-th continuous differentiable functions on A, respectively. We denote $L^\infty(A)$ by the set of bounded measurable functions and $\|\cdot\|_{L^\infty(A)}$ denotes the superemum norm. Let $\mathbb{T}^n$ denote the n-dimensional torus and we identify $\mathbb{T}^n$ with $[0,1]^n$. Define $\Pi : \mathbb{R}^n \to \mathbb{T}^n$ as the canonical projection. By abuse of notations, we denote the periodic extensions of any set $B \subset \mathbb{T}^n$ and any function $f \in C(\mathbb{T}^n)$ to the whole space $\mathbb{R}^n$ by B, and f themselves respectively. For $a, b \in \mathbb{R}$, we write $a \wedge b = \min\{a, b\}$ and $a \vee b = \max\{a, b\}$. We call a function $m : [0,\infty) \to [0,\infty)$ a *modulus* if it is continuous, nondecreasing on $[0,\infty)$ and $m(0) = 0$.

2. Homogenization Results

Lemma 2.1 (Barrier Functions)**.** *We define the functions $\varphi_i^\pm : \mathbb{R}^n \times [0,T] \to \mathbb{R}$ by*

$$\begin{cases} \varphi_1^\pm(x,t) = \dfrac{f_1(x)+f_2(x)}{2} + \dfrac{f_1(x)-f_2(x)}{2} e^{-\frac{2t}{\varepsilon}} \pm Ct \\ \varphi_2^\pm(x,t) = \dfrac{f_1(x)+f_2(x)}{2} + \dfrac{f_2(x)-f_1(x)}{2} e^{-\frac{2t}{\varepsilon}} \pm Ct. \end{cases} \tag{2.1}$$

If we choose $C \geq \max_{i=1,2} \max_{(\xi,p)\in\mathbb{R}^n\times B(0,r)} |H_i(\xi,p)|$, where $r = \|Df_1\|_{L^\infty(\mathbb{R}^n)} + \|Df_2\|_{L^\infty(\mathbb{R}^n)}$, then $(\varphi_1^-, \varphi_2^-)$ and $(\varphi_1^+, \varphi_2^+)$ are, respectively, a subsolution and a supersolution of (C_ε), and

$$(\varphi_1^-, \varphi_2^-)(\cdot,0) = (\varphi_1^+, \varphi_2^+)(\cdot,0) = (f_1, f_2) \text{ on } \mathbb{R}^n.$$

In particular, $\varphi_i^- \leq u_i^\varepsilon \leq \varphi_i^+$ on $\mathbb{R}^n \times [0,T]$ for $i = 1, 2$.

Proof. We calculate that

$$\begin{aligned} &(\varphi_1^-)_t + H_1(\frac{x}{\varepsilon}, D\varphi_1^-) + \frac{1}{\varepsilon}(\varphi_1^- - \varphi_2^-) \\ =& -\frac{f_1(x)-f_2(x)}{\varepsilon} e^{-\frac{2t}{\varepsilon}} - C + H_1(\frac{x}{\varepsilon}, D\varphi_1^-) + \frac{f_1(x)-f_2(x)}{\varepsilon} e^{-\frac{2t}{\varepsilon}} \\ =& -C + H_1(\frac{x}{\varepsilon}, D\varphi_1^-) \leq 0 \end{aligned}$$

for $C > 0$ large enough as chosen above. Similar calculations give us that $(\varphi_1^-, \varphi_2^-)$ and $(\varphi_1^+, \varphi_2^+)$ are, respectively, a subsolution and a supersolution of (C_ε). By the comparison principle for (C_ε) (see [15, 23]) we get $\varphi_i^- \le u_i^\varepsilon \le \varphi_i^+$ on $\mathbb{R}^n \times [0,T]$ for $i = 1, 2$. □

Proof of Theorem 1.2. By Lemma 2.1 we can take the following half-relaxed limits

$$\begin{cases} W(x,t) := \limsup^*_{\varepsilon\to 0} \sup_{i=1,2}[u_i^\varepsilon](x,t) \\ w(x,t) := \liminf_{*\varepsilon\to 0} \inf_{i=1,2}[u_i^\varepsilon](x,t). \end{cases}$$

We now show that W and w are, respectively, a subsolution and a supersolution of (1.3) in $\mathbb{R}^n \times (0,T)$ by employing the perturbed test function method.

Since we can easily check $W(\cdot,0) = w(\cdot,0) = \overline{f}$ on $\mathbb{R}^n$ due to Lemma 2.1, it is enough to prove that W and w are a subsolution and a supersolution, respectively, of the equation in (1.3). We only prove that W is a subsolution since by symmetry we can prove that w is a supersolution. We take a test function $\phi \in C^1(\mathbb{R}^n \times (0,T))$ such that $W - \phi$ has a strict maximum at $(x_0, t_0) \in \mathbb{R}^n \times (0,T)$. Let $P := D\phi(x_0, t_0)$. Choose a sequence $\varepsilon_m \to 0$ such that

$$W(x_0,t_0) = \limsup^*_{m\to\infty} \max_{i=1,2} u_i^{\varepsilon_m}(x_0,t_0).$$

We define the perturbed test functions $\psi_i^{\varepsilon,\alpha}$ for $i = 1, 2$ and $\alpha > 0$ by

$$\psi_i^{\varepsilon,\alpha}(x,y,t) := \phi(x,t) + \varepsilon v_i(\frac{y}{\varepsilon}) + \frac{|x-y|^2}{2\alpha^2},$$

where (v_1, v_2) is a solution of (E_P). By the usual argument in the theory of viscosity solutions, for every $m \in \mathbb{N}$, $\alpha > 0$, there exist $i_{m,\alpha} \in \{1,2\}$ and $(x_{m,\alpha}, y_{m,\alpha}, t_{m,\alpha}) \in \mathbb{R}^n \times \mathbb{R}^n \times (0,T)$ such that

$$\max_{i=1,2} \max_{\mathbb{R}^n\times\mathbb{R}^n\times[0,T]} [u_i^{\varepsilon_m}(x,t) - \psi_i^{\varepsilon_m,\alpha}(x,y,t)] = u_{i_{m,\alpha}}^{\varepsilon_m}(x_{m,\alpha},t_{m,\alpha}) - \psi_{i_{m,\alpha}}^{\varepsilon_m,\alpha}(x_{m,\alpha},y_{m,\alpha},t_{m,\alpha}) \quad (2.2)$$

and up to passing some subsequences

$$\begin{aligned} &(x_{m,\alpha}, y_{m,\alpha}, t_{m,\alpha}) \to (x_m, x_m, t_m) \text{ as } \alpha \to 0, \\ &i_{m,\alpha} \to i_m \in \{1,2\} \text{ as } \alpha \to 0, \\ &(x_m, t_m) \to (x_0, t_0) \text{ as } m \to \infty, \\ &\lim_{m\to\infty} \lim_{\alpha\to 0} u_{i_{m,\alpha}}^{\varepsilon_m}(x_{m,\alpha},t_{m,\alpha}) = W(x_0,t_0). \end{aligned}$$

Choose $j_{m,\alpha}, j_m \in \{1,2\}$ such that $\{i_{m,\alpha}, j_{m,\alpha}\} = \{i_m, j_m\} = \{1,2\}$. By the definition of viscosity solutions, we have

$$\phi_t(x_{m,\alpha},t_{m,\alpha}) + H_{i_{m,\alpha}}(\frac{x_{m,\alpha}}{\varepsilon_m}, D\phi(x_{m,\alpha},t_{m,\alpha}) + \frac{x_{m,\alpha}-y_{m,\alpha}}{\alpha^2}) + \frac{1}{\varepsilon_m}(u_{i_{m,\alpha}}^{\varepsilon_m} - u_{j_{m,\alpha}}^{\varepsilon_m})(x_{m,\alpha},t_{m,\alpha}) \le 0. \quad (2.3)$$

Since (v_1, v_2) is a supersolution of (E_P), we have

$$H_{i_{m,\alpha}}(\frac{y_{m,\alpha}}{\varepsilon_m}, P + \frac{x_{m,\alpha} - y_{m,\alpha}}{\alpha^2}) + (v_{i_{m,\alpha}} - v_{j_{m,\alpha}})(\frac{y_{m,\alpha}}{\varepsilon_m}) \geq \overline{H}(P). \tag{2.4}$$

Let $\alpha \to 0$ in (2.3) and (2.4) to derive

$$\phi_t(x_m, t_m) + H_{i_m}(\frac{x_m}{\varepsilon_m}, D\phi(x_m, t_m) + Q_m) + \frac{1}{\varepsilon_m}(u_{i_m}^{\varepsilon_m} - u_{j_m}^{\varepsilon_m})(x_m, t_m) \leq 0 \tag{2.5}$$

and

$$H_{i_m}(\frac{x_m}{\varepsilon_m}, P + Q_m) + (v_{i_m} - v_{j_m})(\frac{x_m}{\varepsilon_m}) \geq \overline{H}(P), \tag{2.6}$$

where $Q_m := \lim_{\alpha\to 0}(x_{m,\alpha} - y_{m,\alpha})/\alpha^2$. Noting that the correctors v_i are Lipschitz continuous due to the coercivity of H_i, we see that $|Q_m| \leq C$ for $C > 0$ which is independent of m. Combine (2.5) with (2.6) to get

$$\begin{aligned}
\phi_t(x_m, t_m) + \overline{H}(P) &\leq H_{i_m}(\frac{x_m}{\varepsilon_m}, P + Q_m) - H_{i_m}(\frac{x_m}{\varepsilon_m}, D\phi(x_m, t_m) + Q_m) \\
&\quad + \frac{1}{\varepsilon_m}[u_{j_m}^{\varepsilon_m}(x_m, t_m) - (\phi(x_m, t_m) + \varepsilon_m v_{j_m}(\frac{x_m}{\varepsilon_m}))] \\
&\quad - \frac{1}{\varepsilon_m}[u_{i_m}^{\varepsilon_m}(x_m, t_m) - (\phi(x_m, t_m) + \varepsilon_m v_{i_m}(\frac{x_m}{\varepsilon_m}))] \\
&\leq \sigma(|P - D\phi(x_m, t_m)|)
\end{aligned}$$

for some modulus σ. Letting $m \to \infty$, we get the result.

We finally prove that u is Lipschitz continuous. We can easily see that $\overline{f} \pm Mt$ are a supersolution and a subsolution of (1.3), respectively, for $M > 0$ large enough. By the comparison principle for (1.3) we have $|u(x,t) - \overline{f}(x)| \leq Mt$ for all $(x,t) \in \mathbb{R}^n \times [0,T]$. Moreover, the comparison principle for (1.3) also yields that

$$\begin{aligned}
&\sup_{x\in\mathbb{R}^n} |u(x, t+s) - u(x,t)| \leq \sup_{x\in\mathbb{R}^n} |u(x,s) - \overline{f}(x)| \leq Ms \text{ for all } t, s \geq 0, \text{ and} \\
&\sup_{x\in\mathbb{R}^n} |u(x+z, t) - u(x,t)| \leq \sup_{x\in\mathbb{R}^n} |\overline{f}(x+z) - \overline{f}(x)| \leq r|z| \text{ for all } z \in \mathbb{R}^n,\ t \geq 0.
\end{aligned}$$

The proof is complete. □

3. Initial layers and Rate of convergence

3.1. **Inner solutions, Outer solutions, and Matched solutions.** We first derive inner solutions, outer solutions and perform the matching asymptotic expansion method to find matched solutions in a heuristic mode.

As we already obtained in Section 2, outer solutions are same as the limit u give in Theorem 1.2. Now we need to find a right scaling for inner solutions. We let

$$w_i^\varepsilon(x,t) = u_i^\varepsilon(x, \frac{t}{\varepsilon}) \text{ for } i = 1, 2,$$

and plug into (C_ε) to obtain

$$(\mathrm{I}_\varepsilon)\qquad \begin{cases} (w_1^\varepsilon)_t + \varepsilon H_1(\dfrac{x}{\varepsilon}, Dw_1^\varepsilon) + (w_1^\varepsilon - w_2^\varepsilon) = 0 & \text{in } \mathbb{R}^n \times (0, T/\varepsilon), \\ (w_2^\varepsilon)_t + \varepsilon H_2(\dfrac{x}{\varepsilon}, Dw_2^\varepsilon) + (w_2^\varepsilon - w_1^\varepsilon) = 0 & \text{in } \mathbb{R}^n \times (T/\varepsilon), \\ w_i^\varepsilon(x,0) = f_i(x) & \text{on } \mathbb{R}^n \text{ for } i = 1,2. \end{cases}$$

We next assume that w_i^ε have the asymptotic expansions of the form

$$w_i^\varepsilon(x,t) = w_i(x,t) + \varepsilon w_{i1}(x,t) + \varepsilon^2 w_{i2}(x,t)\cdots, \text{ for } i = 1,2.$$

It is then relatively straightforward to see that (w_1, w_2) solves

$$(\mathrm{I})\qquad \begin{cases} (w_1)_t + (w_1 - w_2) = 0 & \text{in } \mathbb{R}^n \times (0,\infty), \\ (w_2)_t + (w_2 - w_1) = 0 & \text{in } \mathbb{R}^n \times (0,\infty), \\ w_i(x,0) = f_i(x) & \text{on } \mathbb{R}^n \text{ for } i = 1,2. \end{cases}$$

Thus, we can compute the explicit formula for the inner solutions

$$\begin{aligned} &(w_1(x,t), w_2(x,t)) \\ &= \left(\frac{f_1(x) + f_2(x)}{2} + \frac{f_1(x) - f_2(x)}{2} e^{-2t},\ \frac{f_1(x) + f_2(x)}{2} + \frac{f_2(x) - f_1(x)}{2} e^{-2t} \right). \end{aligned}$$

The final step is to obtain the matched solutions. We have in this particular situation

$$\lim_{t\to 0} u(x,t) = \lim_{t\to\infty} w_i(x,t) = \frac{f_1(x) + f_2(x)}{2},$$

which shows that the common part of the inner and outer solutions is $(f_1 + f_2)(x)/2$. Hence, the matched solutions are

$$\begin{aligned} &\left(u(x,t) + w_1(x, \frac{t}{\varepsilon}) - \frac{f_1(x) + f_2(x)}{2}, u(x,t) + w_2(x, \frac{t}{\varepsilon}) - \frac{f_1(x) + f_2(x)}{2} \right) \\ &= \left(u(x,t) + \frac{f_1(x) - f_2(x)}{2} e^{-\frac{2t}{\varepsilon}}, u(x,t) + \frac{f_2(x) - f_1(x)}{2} e^{-\frac{2t}{\varepsilon}} \right) \\ &= (m_1^\varepsilon(x,t), m_2^\varepsilon(x,t)), \end{aligned}$$

where m_i^ε are the functions defined by (1.4).

As we can see, the matched solutions contain the layer parts which are essentially the same like the subsolutions and supersolutions that we build in Lemma 2.1. For any fixed $t > 0$, we can see that $(m_1^\varepsilon(x,t), m_2^\varepsilon(x,t))$ converges to $(u(x,t), u(x,t))$ exponentially fast. But for $t = O(\varepsilon)$ then we do not have such convergence. In particular, we have $(m_1^\varepsilon(x,\varepsilon), m_2^\varepsilon(x,\varepsilon))$ converges to $(u(x,t) + (f_1 - f_2)(x)/(2e^2), u(x,t) + (f_2 - f_1)(x)/(2e^2))$. On the other hand, the fact that $((m_1^\varepsilon)_t, (m_2^\varepsilon)_t)$ is not bounded also give us an intuition about the unboundedness of $((u_1^\varepsilon)_t, (u_2^\varepsilon)_t)$.

It is therefore interesting if we can study the behavior of the difference between the real solutions $(u_1^\varepsilon, u_2^\varepsilon)$ and the matched solutions $(m_1^\varepsilon, m_2^\varepsilon)$.

3.2. **Rate of convergence to matched solutions.** In this subsection, we assume further that

(A3) H_i are (uniformly) Lipschitz in the p-variable for $i = 1, 2$, i.e. there exists a constant $C_H > 0$ such that

$$|H_i(\xi, p) - H_i(\xi, q)| \le C_H|p - q| \text{ for all } \xi \in \mathbb{T}^n \text{ and } p, q \in \mathbb{R}^n.$$

We now prove Theorem 1.3 by splitting $\mathbb{R}^n \times [0, T]$ into two parts, which are $\mathbb{R}^n \times [0, \varepsilon|\log\varepsilon|]$ and $\mathbb{R}^n \times [\varepsilon|\log\varepsilon|, T]$. For the part of small time $\mathbb{R}^n \times [0, \varepsilon|\log\varepsilon|]$, we use the barrier functions in Lemma 2.1 and the effective equation to obtain the results. The L^∞-bounds of $|u_i^\varepsilon - m_i^\varepsilon|$ for $i = 1, 2$ on $\mathbb{R}^n \times [\varepsilon|\log\varepsilon|, T]$ can be obtained by using techniques similar to those of Capuzzo-Dolcetta and Ishii [9].

Proposition 3.1 (Initial Layer)**.** *There exists $C > 0$ such that*

$$|(u_i^\varepsilon - m_i^\varepsilon)(x, t)| \le C\varepsilon|\log\varepsilon| \text{ for all } (x, t) \in \mathbb{R}^n \times [0, \varepsilon|\log\varepsilon|] \text{ and } i = 1, 2.$$

Proof. We only prove the case $i = 1$. By symmetry we can prove the case $i = 2$. Let $C \ge \max_{i=1,2} \max_{(\xi,p)\in\mathbb{R}^n\times B(0,r)} |H_i(\xi, p)|$ be a constant, where $r = \|Df_1\|_{L^\infty(\mathbb{R}^n)} + \|Df_2\|_{L^\infty(\mathbb{R}^n)}$ and note that u is Lipschitz continuous with a Lipschitz constant $C_u := (1/2)(\|Df_1\|_\infty + \|Df_2\|_\infty)$. By Lemma 2.1 we have

$$\begin{aligned}
&\big|u_1^\varepsilon(x, t) - (u(x, t) + \frac{f_1(x) - f_2(x)}{2} e^{-\frac{2t}{\varepsilon}})\big| \\
&\le \big|u(x, t) - \frac{f_1(x) + f_2(x)}{2}\big| + Ct = |u(x, t) - u(x, 0)| + Ct \\
&\le (C + C_u)t \le (C + C_u)\varepsilon|\log\varepsilon|
\end{aligned}$$

for all $t \in [0, \varepsilon|\log\varepsilon|]$. □

Proposition 3.2. *Assume that* (A3) *holds. For $T > 0$ there exists $C = C(T) > 0$ such that*

$$|u_i^\varepsilon(x, t) - u(x, t)| \le C\varepsilon^{1/3} \quad \text{for } (x, t) \in \mathbb{R}^n \times [\varepsilon|\log\varepsilon|, T] \text{ and } i = 1, 2.$$

Lemma 3.3. *Assume that* (A3) *holds. For each $\delta > 0$ and $P \in \mathbb{R}^n$, there exists a unique solution $(v_1^\delta, v_2^\delta) \in C^{0,1}(\mathbb{T}^n)^2$ of*

$$(\mathrm{E}_P^\delta) \qquad \begin{cases} H_1(\xi, P + Dv_1^\delta(\xi, P)) + (1 + \delta)v_1^\delta(\xi, P) - v_2^\delta(\xi, P) = 0 & \text{in } \mathbb{T}^n, \\ H_2(\xi, P + Dv_2^\delta(\xi, P)) + (1 + \delta)v_2^\delta(\xi, P) - v_1^\delta(\xi, P) = 0 & \text{in } \mathbb{T}^n. \end{cases}$$

Moreover,

(i) *there exists a constant $C > 0$ independent of δ such that*

$$\delta|v_i^\delta(\xi, P) - v_i^\delta(\xi, Q)| \le C|P - Q| \text{ for } \xi \in \mathbb{T}^n, P, Q \in \mathbb{R}^n \text{ and } i = 1, 2;$$

(ii) *for each $R > 0$, there exists a constant $C = C(R) > 0$ independent of δ such that*

$$|\delta v_i^\delta(\xi, P) + \overline{H}(P)| \le C(R)\delta \text{ for } \xi \in \mathbb{T}^n, P \in \overline{B}(0, R) \text{ and } i = 1, 2.$$

Proof. By the classical result (see [15, 23]) we can easily see that there exists a unique solution (v_1^δ, v_2^δ) of (E_P^δ) for any $P \in \mathbb{R}^n$. We see that $\big(v_1^\delta(\cdot, P) \pm C_H|P - Q|/\delta, v_2^\delta(\cdot, P) \pm C_H|P - Q|/\delta\big)$ are a supersolution and subsolution of (E_Q^δ), respectively, in view of (A3). By the comparison principle for (E_Q^δ) we have

$$v_i^\delta(\xi, P) - \frac{C_H|P - Q|}{\delta} \le v_i^\delta(\xi, Q) \le v_i^\delta(\xi, P) + \frac{C_H|P - Q|}{\delta} \text{ for } \xi \in \mathbb{T}^n,\ i = 1, 2,$$

which completes (i).

Let $C_1(P) = \max_{i=1,2} \max_{\xi \in \mathbb{T}^n} |H_i(\xi, P)|$. It is clear that $(-C_1(P)/\delta, -C_1(P)/\delta)$ and $(C_1(P)/\delta, C_1(P)/\delta)$ are a subsolution and a supersolution of (E_P^δ), respectively. Note that

$$|H_i(\xi, P)| \le |H_i(\xi, 0)| + |H_i(\xi, 0) - H_i(\xi, P)| \le C(1 + |P|)$$

for $C \ge \max_{i=1,2,\, \xi \in \mathbb{T}^n} |H_i(\xi, 0)| \vee C_H$. Therefore, by the comparison principle again we get

$$\delta \|v_i^\delta(\cdot, P)\|_{L^\infty(\mathbb{T}^n)} \le C_1(P) \le C(1 + |P|). \tag{3.1}$$

Next, sum up the two equations of (E_P^δ) to get

$$H_1(\xi, P + Dv_1^\delta(\xi, P)) + H_2(\xi, P + Dv_2^\delta(\xi, P)) \le 2C(1 + |P|).$$

Thus, for each $R > 0$, there exists a constant $C = C(R) \ge 0$ so that

$$\|Dv_i^\delta(\cdot, P)\|_{L^\infty(\mathbb{T}^n)} \le C(R) \text{ for } |P| \le R \text{ and } i = 1, 2. \tag{3.2}$$

We look back at (E_P^δ) and take the inequalities (3.1), (3.2) into account to deduce that

$$\|v_1^\delta(\cdot, P) - v_2^\delta(\cdot, P)\|_{L^\infty(\mathbb{T}^n)} \le C(R) \text{ for } |P| \le R. \tag{3.3}$$

Let $\mu^+ := \max_{i=1,2,\, \xi \in \mathbb{T}^n} \delta v_i^\delta(\xi, P)$ and $\mu^- := \min_{i=1,2,\, \xi \in \mathbb{T}^n} \delta v_i^\delta(\xi, P)$. Then we have

$$\mu^- \le -\overline{H}(P) \le \mu^+. \tag{3.4}$$

Indeed, suppose that $\mu^+ < -\overline{H}(P)$ and then by the comparison principle we have $v_i^\delta \ge w_i$ on $\mathbb{T}^n$ for any solution $(w_1, w_2, \overline{H}(P))$ of (E_P). This is a contradiction, since for any $C_2 \in \mathbb{R}$ $(w_1 + C_2, w_2 + C_2, \overline{H}(P))$ is a solution of (E_P) too. Similarly we see that $\mu^- \le -\overline{H}(P)$. By (3.2)–(3.4) we can get the desired conclusion of (ii). □

Proof of Proposition 3.2. Let $(v_1^\delta(\cdot,P), v_2^\delta(\cdot,P))$ be the solution of (E_P^δ) for $P \in \mathbb{R}^n$. We consider the auxiliary functions

$$\Phi_i(x,y,t,s) := u_i^\varepsilon(x,t) - u(y,s) - \varepsilon v_i^\delta(\frac{x}{\varepsilon}, \frac{x-y}{\varepsilon^\beta}) - \frac{|x-y|^2+(t-s)^2}{2\varepsilon^\beta} - K(t+s)$$

for $i = 1, 2$, where $\delta = \varepsilon^\theta$ and $\beta, \theta \in (0,1)$ and $K \geq 0$ to be fixed later.

For simplicity of explanation we assume that Φ_i takes a global maximum on $\mathbb{R}^{2n} \times [\varepsilon|\log\varepsilon|, T]^2$ and let $(\hat{x}, \hat{y}, \hat{t}, \hat{s})$ be a point such that

$$\max_{i=1,2} \max_{\mathbb{R}^n \times [\varepsilon|\log\varepsilon|, T]} \Phi_i(x,y,t,s) = \Phi_1(\hat{x}, \hat{y}, \hat{t}, \hat{s}). \tag{3.5}$$

For a more rigorous proof we need to add the term $-\gamma|x|^2$ to Ψ_i for $\gamma > 0$. See the proof of Theorem 1.1 in [9] for the detail. We first consider the case where $\hat{t}, \hat{s} > \varepsilon|\log\varepsilon|$.

Claim. If $0 < \theta < 1 - \beta$, then there exists $M > 0$ such that $(|\hat{x} - \hat{y}| + |\hat{t} - \hat{s}|)/\varepsilon^\beta \leq M$.

We use $\Phi_1(\hat{x}, \hat{y}, \hat{t}, \hat{s}) \geq \Phi_1(\hat{x}, \hat{x}, \hat{t}, \hat{t})$, Lemma 3.3 (i) and that u is Lipschitz continuous to deduce that

$$\begin{aligned}
\frac{|\hat{x}-\hat{y}|^2 + |\hat{t}-\hat{s}|^2}{2\varepsilon^\beta} &\leq |u(\hat{x},\hat{t}) - u(\hat{y},\hat{s})| + \varepsilon|v_1^\delta(\frac{\hat{x}}{\varepsilon}, \frac{\hat{x}-\hat{y}}{\varepsilon^\beta}) - v_1^\delta(\frac{\hat{x}}{\varepsilon}, 0)| + K|\hat{t}-\hat{s}| \\
&\leq C_u(|\hat{x}-\hat{y}| + |\hat{t}-\hat{s}|) + C\varepsilon\frac{1}{\varepsilon^\theta}\frac{|\hat{x}-\hat{y}|}{\varepsilon^\beta} + K|\hat{t}-\hat{s}| \\
&\leq C'(|\hat{x}-\hat{y}| + |\hat{t}-\hat{s}|)
\end{aligned}$$

for some $C, C' > 0$, which implies the result of Claim.

We fix $(y,s) = (\hat{y}, \hat{s})$ and notice that the function

$$(x,t) \mapsto u_1^\varepsilon(x,t) - \varepsilon v_i^\delta(\frac{x}{\varepsilon}, \frac{x-\hat{y}}{\varepsilon^\beta}) - \frac{|x-\hat{y}|^2 + (t-\hat{s})^2}{2\varepsilon^\beta} - Kt$$

attains the maximum at $(\hat{x}, \hat{t})$. For $\alpha > 0$, we define the function ψ by

$$\psi(x,\xi,z,t) := u_1^\varepsilon(x,t) - \varepsilon v_1^\delta(\xi, \frac{z-\hat{y}}{\varepsilon^\beta}) - \frac{|x-\hat{y}|^2 + |t-\hat{s}|^2}{2\varepsilon^\beta} - \frac{|x-\varepsilon\xi|^2 + |x-z|^2}{2\alpha} - Kt.$$

Let ψ attain the maximum at $(x_\alpha, \xi_\alpha, z_\alpha, t_\alpha)$ and then we may assume that $(x_\alpha, \xi_\alpha, z_\alpha, t_\alpha) \to (\hat{x}, \hat{x}/\varepsilon, \hat{x}, \hat{t})$ as $\alpha \to 0$ up to passing some subsequences if necessary. By the definition of viscosity solutions, we have

$$K + \frac{t_\alpha - \hat{s}}{\varepsilon^\beta} + H_1(\frac{x_\alpha}{\varepsilon}, \frac{x_\alpha - \hat{y}}{\varepsilon^\beta} + \frac{x_\alpha - \varepsilon\xi_\alpha}{\alpha} + \frac{x_\alpha - z_\alpha}{\alpha}) + \frac{1}{\varepsilon}(u_1^\varepsilon - u_2^\varepsilon)(x_\alpha, t_\alpha) \leq 0, \tag{3.6}$$

and

$$H_1(\xi_\alpha, \frac{z_\alpha - \hat{y}}{\varepsilon^\beta} + \frac{x_\alpha - \varepsilon\xi_\alpha}{\alpha}) + (1+\delta)v_1^\delta(\xi_\alpha, \frac{z_\alpha - \hat{y}}{\varepsilon^\beta}) - v_2^\delta(\xi_\alpha, \frac{z_\alpha - \hat{y}}{\varepsilon^\beta}) \geq 0. \tag{3.7}$$

Next, since $\psi(x_\alpha, \xi_\alpha, z_\alpha, t_\alpha) \geq \psi(x_\alpha, \xi_\alpha, x_\alpha, t_\alpha)$ we get

$$\frac{|x_\alpha - z_\alpha|^2}{2\alpha} \leq \varepsilon(v_1^\delta(\xi_\alpha, \frac{x_\alpha - \hat{y}}{\varepsilon^\beta}) - v_1^\delta(\xi_\alpha, \frac{z_\alpha - \hat{y}}{\varepsilon^\beta})) \leq C\varepsilon^{1-\theta-\beta}|x_\alpha - z_\alpha|$$

by Lemma 3.3 (i). Thus, $|x_\alpha - z_\alpha|/\alpha \leq C\varepsilon^{1-\theta-\beta}$. Send $\alpha \to 0$ and we get

$$K + \frac{\hat{t} - \hat{s}}{\varepsilon^\beta} + \overline{H}(\frac{\hat{x} - \hat{y}}{\varepsilon^\beta}) + \frac{1}{\varepsilon}(u_1^\varepsilon - u_2^\varepsilon)(\hat{x}, \hat{t}) - v_1^\delta(\frac{\hat{x}}{\varepsilon}, \frac{\hat{x} - \hat{y}}{\varepsilon^\beta}) + v_2^\delta(\frac{\hat{x}}{\varepsilon}, \frac{\hat{x} - \hat{y}}{\varepsilon^\beta}) - C(\varepsilon^\theta + \varepsilon^{1-\theta-\beta}) \leq 0. \tag{3.8}$$

Similarly we fix $(x, t) = (\hat{x}, \hat{t})$ and do a similar procedure to the above to obtain

$$-K + \frac{\hat{t} - \hat{s}}{\varepsilon^\beta} + \overline{H}(\frac{\hat{x} - \hat{y}}{\varepsilon^\beta}) + C(\varepsilon^\theta + \varepsilon^{1-\theta-\beta}) \geq 0. \tag{3.9}$$

Combining (3.8), (3.9), and (3.5), we get

$$2K \leq C(\varepsilon^\theta + \varepsilon^{1-\theta-\beta}). \tag{3.10}$$

Now we choose $\theta = \beta = 1/3$ and $K = K_1\varepsilon^{1/3}$ for K_1 large enough to get the contradiction in (3.10). Hence either $\hat{t} = -\varepsilon \log \varepsilon$ or $\hat{s} = -\varepsilon \log \varepsilon$ holds. The proof is complete immediately. □

Theorem 1.3 is a straightforward result of Propositions 3.1, 3.2.

4. Properties of effective Hamiltonians

4.1. **Elementary properties.**

Proposition 4.1.
(i) (*Coercivity*) $\overline{H}(P) \to +\infty$ *as* $|P| \to \infty$.
(ii) (*Convexity*) *If* H_i *are convex in the p-variable for* $i = 1, 2$, *then* $\overline{H}$ *is convex.*

Proof. (i) For each $\delta > 0$ and $P \in \mathbb{R}^n$, let (v_1^δ, v_2^δ) be a solution of (E_P^δ) and without loss of generality, we may assume that $v_1^\delta(\xi_0, P) = \max_{i=1,2,\,\xi\in\mathbb{T}^n} v_i^\delta(\xi, P)$ for some $\xi_0 \in \mathbb{T}^n$. By the definition of viscosity solutions we have $H_1(\xi_0, P) \leq H_1(\xi_0, P) + (v_1^\delta - v_2^\delta)(\xi_0, P) \leq -\delta v_1^\delta(\xi_0, P)$. We let $\delta \to 0$ to derive that $\overline{H}(P) \geq \min_{i=1,2,\,\xi\in\mathbb{T}^n} H_i(\xi, P)$. Since H_i are coercive for $i = 1, 2$, so is $\overline{H}$.

(ii) We argue by contradiction. Suppose that $\overline{H}$ is not convex and then there would exist $P, Q \in \mathbb{R}^n$ such that

$$2\varepsilon_0 := \overline{H}(\frac{P + Q}{2}) - \frac{\overline{H}(P) + \overline{H}(Q)}{2} > 0. \tag{4.1}$$

We define the functions $w_i \in C(\mathbb{T}^n)$ so that $w_i(\xi) := (v_i(\xi, P) + v_i(\xi, Q))/2$ for $i = 1, 2$, where $(v_1(\cdot, P), v_2(\cdot, P))$ and $(v_1(\cdot, Q), v_2(\cdot, Q))$ are solutions of (E_P) and (E_Q), respectively.

Due to the convexity of H_i for $i = 1, 2$ we have

$$\begin{cases} H_1(\xi, \dfrac{P+Q}{2} + Dw_1(\xi)) + w_1(\xi) - w_2(\xi) \le \dfrac{\overline{H}(P) + \overline{H}(Q)}{2}, \\ H_2(x, \dfrac{P+Q}{2} + Dw_2(\xi)) + w_2(\xi) - w_1(\xi) \le \dfrac{\overline{H}(P) + \overline{H}(Q)}{2}. \end{cases}$$

By (4.1) there exists a small constant $\delta > 0$ such that

$$\begin{cases} H_1(\xi, \dfrac{P+Q}{2} + Dw_1(\xi)) + (1+\delta)w_1(\xi) - w_2(\xi) \le \overline{H}(\dfrac{P+Q}{2}) - \varepsilon_0, \\ H_2(x, \dfrac{P+Q}{2} + Dw_2(\xi)) + (1+\delta)w_2(\xi) - w_1(\xi) \le \overline{H}(\dfrac{P+Q}{2}) - \varepsilon_0, \end{cases}$$

and

$$\begin{cases} H_1(\xi, \dfrac{P+Q}{2} + Dv_1(\xi, \dfrac{P+Q}{2})) + (1+\delta)v_1(\xi, \dfrac{P+Q}{2}) - v_2(\xi, \dfrac{P+Q}{2}) \ge \overline{H}(\dfrac{P+Q}{2}) - \varepsilon_0, \\ H_2(\xi, \dfrac{P+Q}{2} + Dv_2(\xi, \dfrac{P+Q}{2})) + (1+\delta)v_2(\xi, \dfrac{P+Q}{2}) - v_1(\xi, \dfrac{P+Q}{2}) \ge \overline{H}(\dfrac{P+Q}{2}) - \varepsilon_0. \end{cases}$$

By the comparison principle we get

$$\frac{v_i(\xi, P) + v_i(\xi, Q)}{2} \le v_i(\xi, \frac{P+Q}{2}) \qquad \text{for } i = 1, 2. \tag{4.2}$$

Notice that (4.2) is still correct even if we replace $v_i(\xi, (P+Q)/2)$ by $v_i(\xi, (P+Q)/2) + C_1$ for $i = 1, 2$ and for any $C_1 \in \mathbb{R}$, which yields the contradiction. □

The uniqueness of the effective Hamiltonian for (E_P) and the cell problem for single Hamilton–Jacobi equations give the following proposition.

Proposition 4.2. *If $H_1 = H_2 = K$, then*

$$\overline{H}(P) = \overline{K}(P) \text{ for all } P \in \mathbb{R}^n,$$

where $\overline{K}$ is the effective Hamiltonian corresponding to K.

Proposition 4.3. *If H_i are homogeneous with degree 1 in the p-variable for $i = 1, 2$, then $\overline{H}$ is positive homogeneous with degree 1.*

Proof. Let $(v_1, v_2, \overline{H}(P))$ be a solution of (E_P) for any $P \in \mathbb{R}^n$. If H_i is homogeneous with degree 1 in the p-variable, then $(rv_1, rv_2, r\overline{H}(P))$ is a solution of (E_{rP}) for any $r > 0$. Therefore by the uniqueness of the effective Hamiltonian we get the conclusion. □

Proposition 4.4. *We define the Hamiltonian K as*

$$K(\xi, p) := \max\{H_1(\xi, p), H_2(\xi, p)\}.$$

Let $\overline{K}$ be its corresponding effective Hamiltonian and then

$$\overline{H}(P) \le \overline{K}(P).$$

Proof. For each $P \in \mathbb{R}^n$, there exists $\varphi(\cdot, P) \in C^{0,1}(\mathbb{T}^n)$ such that

$$K(\xi, P + D\varphi(\xi, P)) = \overline{K}(P).$$

Thus $(\varphi(\cdot, P), \varphi(\cdot, P), \overline{K}(P))$ is a subsolution of (E_P). We hence get $\overline{K}(P) \geq \overline{H}(P)$ by Proposition 4.6. □

We give an example that we can calculate the effective Hamiltonian explicitly.

Example 4.5. Let $n = 1$ and $H_1(\xi, p) = |p|$, $H_2(\xi, p) = a(\xi)|p|$, where

$$a(\xi) := \frac{1 - (\frac{1}{8\pi^2}\cos(2\pi\xi) + \frac{1}{4\pi}\sin(2\pi\xi))}{1 + (\frac{1}{2} + \frac{1}{8\pi^2})\cos(2\pi\xi)} > 0.$$

By Proposition 4.3 we have $\overline{H}(P) = \overline{H}(1)P$ for $P \geq 0$. Set

$$v_1(\xi, 1) := \frac{1}{16\pi^3}\sin(2\pi\xi) - \frac{1}{8\pi^2}\cos(2\pi\xi),\ v_2(\xi, 1) := (\frac{1}{4\pi} + \frac{1}{16\pi^3})\sin(2\pi\xi).$$

Then we can confirm that $(v_1(\cdot, 1), v_2(\cdot, 1), 1)$ is a solution of (E_1). Therefore $\overline{H}(1) = 1$ and thus, $\overline{H}(P) = P$ for $P \geq 0$.

For any $P < 0$ we have $\overline{H}(P) = \overline{H}(-1) \cdot (-P)$. Set

$$v_1(\xi, -1) := -\big(\frac{1}{16\pi^3}\sin(2\pi\xi) - \frac{1}{8\pi^2}\cos(2\pi\xi)\big),\ v_2(\xi, 1) := -(\frac{1}{4\pi} + \frac{1}{16\pi^3})\sin(2\pi\xi).$$

Then we can confirm that $(v_1(\cdot, -1), v_2(\cdot, -1), 1)$ is a solution of (E_{-1}). Therefore $\overline{H}(-1) = 1$ and thus, $\overline{H}(P) = -P$ for $P \leq 0$. Thus, we get $\overline{H}(P) = |P|$.

4.2. **Representation formulas for the effective Hamiltonian.** In this subsection we derive representation formulas for the effective Hamiltonian $\overline{H}(P)$. See [10, 20] for the min-max formulas for the effective Hamiltonian for single equations.

Proposition 4.6 (Representation formula 1)**.** *We have*

$$\overline{H}(P) = \inf\{c : \text{ there exists } (\phi_1, \phi_2) \in C(\mathbb{T}^n)^2 \text{ so that the triplet } (\phi_1, \phi_2, c) \text{ is a subsolution of } (\mathrm{E_P})\}. \quad (4.3)$$

Proof. Fix $P \in \mathbb{R}^n$ and we denote by $c(P)$ the right-hand side of (4.3). By the definition of $c(P)$ we can easily see that $\overline{H}(P) \geq c(P)$. We prove the other way around. Assume by contradiction that there exist a triplet $(\phi_1, \phi_2, c) \in C(\mathbb{T}^n)^2 \times \mathbb{R}$ which is a subsolution of (E_P) and $c < \overline{H}(P)$. Let $(v_1, v_2, \overline{H}(P))$ be a solution of (E_P) and take $C > 0$ so that $\phi_i > v_i - C =: \overline{v}_i$ on $\mathbb{T}^n$. Then since $\overline{v}_i$ and ϕ_i are bounded on $\mathbb{T}^n$, for $\varepsilon > 0$ small enough, we have

$$\begin{cases} H_1(\xi, P + D\overline{v}_1) + (1+\varepsilon)\overline{v}_1 - \overline{v}_2 \geq H_1(\xi, P + D\phi_1) + (1+\varepsilon)\phi_1 - \phi_2 \\ H_2(\xi, P + D\overline{v}_2) + (1+\varepsilon)\overline{v}_2 - \overline{v}_1 \geq H_2(\xi, P + D\phi_2) + (1+\varepsilon)\phi_2 - \phi_1. \end{cases}$$

By the comparison principle (see [15, 23]) we deduce $\overline{v}_i \geq \phi_i$ on $\mathbb{T}^n$ which yields the contradiction. □

If we assume the convexity on $H_i(\xi, \cdot)$ for any $\xi \in \mathbb{R}^n$, by the classical result on the representation formula for the effective Hamiltonian for single Hamilton–Jacobi equations we can easily see that

$$\begin{aligned}\overline{H}(P) &= \inf_{\varphi \in C^1(\mathbb{T}^n)} \max_{\xi \in \mathbb{T}^n} [H_1(\xi, P + D\varphi(\xi)) + v_1(\xi, P) - v_2(\xi, P)] \\ &= \inf_{\psi \in C^1(\mathbb{T}^n)} \max_{\xi \in \mathbb{T}^n} [H_2(\xi, P + D\psi(\xi)) + v_2(\xi, P) - v_1(\xi, P)]\end{aligned} \tag{4.4}$$

for any solution $(v_1(\cdot, P), v_2(\cdot, P))$ of (E_P), which is in a sense an implicit formula. For the weakly coupled system we have the following representation formula.

Proposition 4.7 (Representation formula 2)**.** *If H_i are convex in the p-variable for $i = 1, 2$, then*

$$\overline{H}(P) = \inf_{(\phi_1, \phi_2) \in C^1(\mathbb{T}^n)^2} \max_{i=1,2,\, \xi \in \mathbb{T}^n} [H_i(\xi, P + D\phi_i(\xi)) + \phi_i(\xi) - \phi_j(\xi)], \tag{4.5}$$

where we take $j \in \{1, 2\}$ so that $\{i, j\} = \{1, 2\}$.

Lemma 4.8. *Assume that H_i are convex in the p-variable. Let $(v_1, v_2, \overline{H}(P)) \in C(\mathbb{T}^n)^2$ be a subsolution of (E_P). Set $v_{i\delta}(x) := \rho_\delta * v_i(x)$, where $\rho_\varepsilon(x) := (1/\varepsilon^n)\rho(x/\varepsilon)$ and $\rho \in C^\infty(\mathbb{R}^n)$ be a standard mollification kernel, i.e., $\rho \geq 0$, $\operatorname{supp}\rho \subset B(0,1)$, and $\int_{\mathbb{R}^n} \rho(x)\,dx = 1$. Then, $(v_{1\delta}, v_{2\delta}, \overline{H}(P) + \omega(\delta))$ is a subsolution of (E_P) for some modulus ω.*

Proof. Note that in view of the coercivity of H_i, v_i are Lipschitz continuous and $(v_1, v_2, \overline{H}(P))$ solves (E_P) almost everywhere. Fix any $\xi \in \mathbb{T}^n$. We calculate that

$$\begin{aligned}\overline{H}(P) &\geq \rho_\delta * \big(H_1(\cdot, Dv_1(\cdot)) + (v_1 - v_2)\big)(\xi) \\ &= \int_{B(\xi,\delta)} \rho_\delta(\xi - \eta)\big(H_1(\eta, Dv_1(\eta)) + (v_1 - v_2)(\eta)\big)\,d\eta \\ &\geq \int_{B(\xi,\delta)} \rho_\delta(\xi - \eta)\big(H_1(\xi, Dv_1(\eta)) - \omega(\delta)\big)\,d\eta + (v_{1\delta} - v_{2\delta})(\xi) \\ &\geq H_1(\xi, \rho_\delta * Dv_1(\xi)) + (v_{1\delta} - v_{2\delta})(\xi) - \omega(\delta) \\ &= H_1(\xi, Dv_{1\delta}(\xi)) + (v_{1\delta} - v_{2\delta})(\xi) - \omega(\delta),\end{aligned}$$

where the third inequality follows by using Jensen's inequality for some modulus ω. □

Proof of Proposition 4.7. Let $c(P)$ be the constant on the right-hand side of (4.5). Noting that for any $(\phi_1, \phi_2) \in C^1(\mathbb{T}^n)^2$

$$H_i(\xi, P + D\phi_i(\xi)) + (\phi_i - \phi_j)(\xi) \leq \max_{i=1,2,\, \xi \in \mathbb{T}^n} [H_i(\xi, P + D\phi_i(\xi)) + (\phi_i - \phi_j)(\xi)] =: a_{\phi_1,\phi_2}$$

for every $\xi \in \mathbb{T}^n$. By Proposition 4.6 we see that $\overline{H}(P) \leq a_{\phi_1,\phi_2}$ for all $(\phi_1, \phi_2) \in C^1(\mathbb{T}^n)^2$. Therefore we get $\overline{H}(P) \leq c(P)$.

Conversely, we observe that by Proposition 1.1 $(v_{1\delta}(\cdot,P), v_{2\delta}(\cdot,P), \overline{H}(P)+\omega(\delta)) \in C^1(\mathbb{T}^n)^2\times\mathbb{R}$ is a subsolution of (E_P). Therefore, by the definition of $c(P)$ we see that $c(P)\le \overline{H}(P)+\omega(\delta)$. Sending $\delta\to 0$ yields the conclusion. □

If H_i are convex in the p-variable, then there is a variational formula of solutions of the initial value problem and the cell problem as stated in Introduction. Therefore, naturally we have the following variational formula

$$\begin{aligned}\overline{H}(P) &= -\lim_{\delta\to 0}\inf_{\eta}\mathbb{E}_i\Big[\int_0^{+\infty} e^{-\delta s}\big(-P\cdot\dot\eta(s)+L_{\nu(s)}(\eta(s),-\dot\eta(s))\big)\,ds\Big]\\ &= -\lim_{t\to\infty}\frac{1}{t}\inf_{\eta}\mathbb{E}_i\Big[\int_0^{t}\big(-P\cdot\dot\eta(s)+L_{\nu(s)}(\eta(s),-\dot\eta(s))\big)\,ds\Big],\end{aligned}$$

where the infimum is taken over $\eta\in \mathrm{AC}([0,+\infty),\mathbb{R}^n)$ such that $\eta(0)=x$ and $\mathbb{E}_i$ denotes the expectation of a process with $\nu(0)=i$ given by (1.2).

Remark 4.9. When we consider the nonconvex Hamilton–Jacobi equations, in general we cannot expect the formula (4.5). Take the Hamiltonian

$$H_i(\xi,p) := (|p|^2-1)^2 \text{ for } i=1,2 \tag{4.6}$$

for instance. In this example if we calculate the right-hand side of (4.5) with $P=0$, then it is 0. But we can easily check that $\overline{H}(0)=1$, since in this case we have solutions (which are constants).

The following formula is a revised min-max formula for the effective Hamiltonian for nonconvex Hamilton–Jacobi equations.

Proposition 4.10. *We have*

$$\overline{H}(P) = \inf_{(\phi_1,\phi_2)\in C^{0,1}(\mathbb{T}^n)^2}\ \max_{i=1,2,\,\xi\in\mathbb{T}^n}\ \sup_{p\in D^+\phi_i(\xi)}[H_i(\xi,P+p)+(\phi_i-\phi_j)(\xi)], \tag{4.7}$$

where if $D^+\phi_i(\xi)=\emptyset$, then we set $\sup_{p\in D^+\phi_i(\xi)}[H_i(\xi,P+p)+(\phi_i-\phi_j)(\xi)] = -\infty$ by convention.

We notice that if H_i are given by (4.6), then the right-hand side of (4.7) with $P=0$ is 1.

Proof. The proof is already in the proof of Proposition 4.7. We just need to be careful for the definition of viscosity subsolutions. Indeed, let c be the right-hand side of (4.7) and noting that for any $(\phi_1,\phi_2)\in C^{0,1}(\mathbb{T}^n)^2$, $\xi\in\mathbb{T}^n$, and $q\in D^+\phi_i(\xi)$,

$$H_i(\xi,P+q)+(\phi_i-\phi_j)(\xi)\le \max_{\xi\in\mathbb{T}^n,i=1,2}\ \sup_{p\in D^+\phi_i(\xi)}[H_i(\xi,P+p)-(\phi_i-\phi_j)(\xi)] =: a_{\phi_1,\phi_2}.$$

Thus, $\overline{H}(P)\le a_{\phi_1,\phi_2}$ for all $(\phi_1,\phi_2)\in C^{0,1}(\mathbb{T}^n)^2$ by Proposition 4.6. Therefore, $\overline{H}(P)\le c$.

Conversely, there exists a viscosity subsolution $(v_1(\cdot,P), v_2(\cdot,P), \overline{H}(P)) \in C^{0,1}(\mathbb{T}^n)^2 \times \mathbb{R}$ of (E_P). By the definition of viscosity subsolutions we have

$$H_i(\xi, P+p) + (v_i - v_j)(\xi) \le \overline{H}(P) \text{ for all } \xi \in \mathbb{T}^n \text{ and } p \in D^+ v_i(\xi).$$

Thus,

$$\max_{\xi\in\mathbb{T}^n, i=1,2} \sup_{p\in D^+v_i(\xi)} [H_i(\xi, P+p) + (v_i - v_j)(\xi)] \le \overline{H}(P),$$

which implies $c \le \overline{H}(P)$. □

4.3. **Flat parts and Non-flat parts near the origin.** In this subsection, we study the results concerning flat parts and non-flat parts of the effective Hamiltonian $\overline{H}$ near the origin. We first point out that there are some cases in which we can obtain similar results to those of Concordel's results for single equations. We present different techniques to obtain these results , namely the min-max formulas, and the construction of subsolutions. In this subsection, we only deal with the Hamiltonians of the form $H_i(\xi, p) = |p|^2 - V_i(\xi)$, where $V_i \in C(\mathbb{T}^n)$ for $i = 1, 2$ unless otherwise stated.

Theorem 4.11. *Assume that $V_i \ge 0$ in $\mathbb{T}^n$ and $\{V_i = 0\} =: U_i \subset \mathbb{T}^n$ for $i = 1, 2$. We assume further that $U_1 \cap U_2 \ne \emptyset$ and there exist open sets W_1, W_2 in $\mathbb{T}^n$, and a vector $q \in \mathbb{R}^n$ such that $\Pi(q + W_2) \Subset (0,1)^n$ and*

$$U_1 \cup U_2 \subset W_1 \subset W_2 \text{ and } \operatorname{dist}(W_1, \partial W_2),\ \operatorname{dist}(U_1 \cup U_2, \partial W_1) > 0, \tag{4.8}$$

then there exists $\gamma > 0$ such that $\overline{H}(P) = 0$ for $|P| \le \gamma$.




Fig. 4.1. The figure of U_i, W_i.




Fig. 4.2. The graph of φ in case $n = 1$.

Proof. Without loss of generality, we may assume that $q = 0$. Take $\xi_0 \in U_1 \cap U_2$. By Proposition 4.7 we have

$$\begin{aligned}\overline{H}(P) &\ge \inf_{(\varphi_1,\varphi_2)\in C^1(\mathbb{T}^n)} \max_{i=1,2} [|P + D\varphi_i(\xi_0)|^2 - V_i(\xi_0) + \varphi_i(\xi_0, P) - \varphi_j(\xi_0, P)] \\ &\ge \inf_{(\varphi_1,\varphi_2)\in C^1(\mathbb{T}^n)} \max_{i=1,2} [\varphi_i(\xi_0, P) - \varphi_j(\xi_0, P)] = 0.\end{aligned}$$

Now, let $d := \min\{\mathrm{dist}(W_1, \partial W_2),\ \mathrm{dist}(U_1 \cup U_2, \partial W_1)\} > 0$. There exists $\varepsilon_0 > 0$ such that

$$V_i(\xi) \geq \varepsilon_0 \quad \text{for } x \in \mathbb{T}^n \setminus W_1,\ i = 1, 2. \tag{4.9}$$

We define a smooth function φ on $\mathbb{T}^n$ such that

$$\begin{aligned} &\varphi(\xi) = -P \cdot \xi \text{ on } W_1,\ \varphi(\xi) = 0 \text{ on } \mathbb{T}^n \setminus W_2, \\ &|D\varphi| \leq \frac{C|P|}{d} \text{ on } \mathbb{T}^n. \end{aligned} \tag{4.10}$$

Notice that

$$|P + D\varphi(\xi)|^2 - V_i(\xi) = \begin{cases} = -V_i(\xi) \leq 0, & \text{on } W_1, \\ \leq \dfrac{C|P|^2}{d^2} - \varepsilon_0, & \text{on } \mathbb{T}^n \setminus W_1. \end{cases}$$

Thus, $|P + D\varphi(\xi)|^2 - V_i(\xi) \leq 0$ on $\mathbb{T}^n$ provided that $|P| \leq d\sqrt{\varepsilon_0}/C =: \gamma$. We hence have that $(\varphi, \varphi, 0)$ is a subsolution of (E_P) for $|P| \leq \gamma$. Therefore $\overline{H}(P) \leq 0$ for $|P| \leq \gamma$ by Proposition 4.6. □

Remark 4.12. (i) In fact, the result of Theorem 4.11 still holds for more general Hamiltonians

$$H_i(\xi, p) := F_i(\xi, p) - V_i(\xi),$$

where $F_i \in C(\mathbb{T}^n \times \mathbb{R}^n)$ and $V_i \in C(\mathbb{T}^n)$ are assumed to satisfy

(a) the functions $p \mapsto F_i(\xi, p)$ are convex and $F_i(\xi, p) \geq F_i(\xi, 0) = 0$ for all $(\xi, p) \in \mathbb{T}^n \times \mathbb{R}^n$,
(b) V_1, V_2 satisfy the conditions of Theorem 4.11.

(ii) Notice that the assumptions $V_i \geq 0$ and $\{V_i = 0\} \neq \emptyset$ for $i = 1, 2$ are just for simplicity. In general, we can normalize V_i by $V_i - \min_{\xi \in \mathbb{T}^n} V_i(\xi)$ to get back to such situation.
(iii) From the proof of Theorem 4.11 we have

$$v_1(\xi, P) = v_2(\xi, P) \quad \text{for all } \xi \in \{V_1 = 0\} \cap \{V_2 = 0\}$$

for any solution $(v_1(\cdot, P), v_2(\cdot, P))$ of (E_P).
(iv) By Proposition 4.4 we can give another proof to Theorem 4.11 as follows. In this case, we explicitly have

$$K(\xi, p) = \max\{|p|^2 - V_1(\xi), |p|^2 - V_2(\xi)\} = |p|^2 - V(\xi)$$

where $V(\xi) = \min\{V_1(\xi), V_2(\xi)\}$. Note that $V \geq 0$ and $\{V = 0\} = \{V_1 = 0\} \cup \{V_2 = 0\}$. Hence, we can either repeat the above proof for single equations to show that $\overline{H}(P) \leq \overline{K}(P) = 0$ for $|P| \leq \gamma$ or we can use Concordel's result directly.

Notice that condition (4.8) is crucial and plays an important role in the construction of the subsolution $(\varphi, \varphi, 0)$ of (E_P) and could not be removed in the proof of Theorem 4.11. We point out in the next Theorem that there are cases when (4.8) does not hold, then the flatness near the origin of $\overline{H}$ does not appear.

Theorem 4.13. *Assume that $V_i \geq 0$ in $\mathbb{T}^n$ and*

$$\{V_1 = 0\} = \{V_2 = 0\} = \{\xi = (\xi_1, \cdots, \xi_n) \in \mathbb{T}^n : \ \xi_j = 1/2 \text{ for } j \geq 2\} =: K. \tag{4.11}$$

The followings hold.

(i) *There exists $\gamma > 0$ such that $\overline{H}(P) = |P_1|^2$ provided that $|P'| \leq \gamma$ for any $P = (P_1, P') \in \mathbb{R} \times \mathbb{R}^{n-1}$.*
(ii) *$\overline{H}(P) \geq |P_1|^2$ for all $P \in \mathbb{R}^n$.*

Proof. Firstly, we prove that $\overline{H}(P) \leq |P_1|^2$ provided that $|P'| \leq \gamma$ for some $\gamma > 0$ small enough by using exactly the same idea in the proof of Theorem 4.11. We build a function $\varphi(\xi) = \varphi(\xi_2, \cdots, \xi_n) \in C^1(\mathbb{T}^n)$, which does not depend on ξ_1, so that

$$\sum_{j=2}^{n} |P_j + \varphi_{\xi_j}(\xi)|^2 - V_i(\xi) \leq 0 \text{ on } \mathbb{T}^n$$

for $i = 1, 2$ and for $|P'| \leq \gamma$ with $\gamma > 0$ small enough. Thus

$$|P + D\varphi(\xi)|^2 - V_i(\xi) \leq |P_1|^2 \text{ on } \mathbb{T}^n$$

for $i = 1, 2$. By Proposition 4.6 $\overline{H}(P) \leq |P_1|^2$.

We now prove that $\overline{H}(P) \geq |P_1|^2$. For each $\xi_0 \in K$, we have in view of (4.4)

$$\begin{aligned}\overline{H}(P) &= \inf_{\varphi \in C^1(\mathbb{T}^n)} \max_{\xi \in \mathbb{T}^n} [|P + D\varphi(\xi)|^2 - V_1(\xi) + v_1(\xi, P) - v_2(\xi, P)] \\ &\geq \inf_{\varphi \in C^1(\mathbb{T}^n)} [|P + D\varphi(\xi_0)|^2 + v_1(\xi_0, P) - v_2(\xi_0, P)],\end{aligned}$$

and similarly

$$\overline{H}(P) \geq \inf_{\psi \in C^1(\mathbb{T}^n)} [|P + D\psi(\xi_0)|^2 + v_2(\xi_0, P) - v_1(\xi_0, P)].$$

Take an arbitrary function $\varphi \in C^1(\mathbb{T}^n)$ and observe that

$$\begin{aligned}&\int_K |P + D\varphi(\xi)|^2 \, d\xi_1 \\ &\geq \int_K |P_1 + \varphi_{\xi_1}(\xi)|^2 \, d\xi_1 = \int_K |P_1|^2 + |\varphi_{\xi_1}(\xi)|^2 + 2P_1\varphi_{\xi_1}(\xi) \, d\xi_1 \\ &\geq \int_K |P_1|^2 + 2P_1\varphi_{\xi_1}(\xi) \, d\xi_1 = |P_1|^2.\end{aligned}$$

Thus, it is clear to see that $\overline{H}(P) \geq |P_1|^2$, which implies the result. □

The above two Theorems describe several examples that we can obtain similar results of the flat part or non-flat part of $\overline{H}$ to those of single Hamilton–Jacobi equations in [11, 12]. Indeed, the structures on the potentials V_i for $i = 1, 2$ are very related in such a way that we obtain the shape of $\overline{H}$ like for single equations. We rely on the idea of building the subsolutions $(\varphi, \psi, \overline{H}(P))$ of (E_P) where $\varphi = \psi$, which does not work in general cases.

Next, we start investigating the properties of $\overline{H}$ in some cases where the structures of the potentials V_i for $i = 1, 2$ are widely different and in general we cannot expect $\overline{H}$ to have simple properties. The next question is that: Can we read of information of the effective Hamiltonian in the case where $\{V_1 = \min_{\xi\in\mathbb{T}^n} V_1(\xi)\} \cap \{V_2 = \min_{\xi\in\mathbb{T}^n} V_2(\xi)\} = \emptyset$?

Theorem 4.14. *Let $n = 1$ and assume that for $\varepsilon_0 > 0$ small enough the following properties hold.*

(a) $\{V_1 = 0\} = [\frac{4}{16}, \frac{12}{16}]$, $\{V_1 = -\varepsilon_0\} = [0, 1] \setminus (\frac{3}{16}, \frac{13}{16})$, *and* $-\varepsilon_0 \le V_1 \le 0$ *on* $\mathbb{T}$ *for some* $\varepsilon_0 > 0$.

(b) $\{V_2 = 0\} = [\frac{7}{16}, \frac{9}{16}]$, $\{V_2 = 2\} = [0, 1] \setminus (\frac{6}{16}, \frac{10}{16})$, *and* $0 \le V_2 \le 2$ *on* $\mathbb{T}$.

There exists $\gamma > 0$ such that $\overline{H}(P) = 0$ for $|P| \le \gamma$.

Lemma 4.15. *We have*

$$\overline{H}(P) \ge -\frac{1}{2} \min_{\xi\in\mathbb{T}^n} (V_1 + V_2)(\xi).$$

Proof. Sum up the two equations in (E_P) to get

$$|P + Dv_1|^2 + |P + Dv_2|^2 - V_1 - V_2 = 2\overline{H}(P),$$

which implies $2\overline{H}(P) \ge -(V_1 + V_2)(\xi))$ for a.e. $\xi \in \mathbb{T}^n$, and the proof is complete. □

Proof of Theorem 4.14. Noting that $\min_{\xi\in\mathbb{T}}(V_1+V_2)(\xi) = 0$, and $\{V_1 = -\varepsilon\}\cap\{V_2 = 0\} = \emptyset$, we have $\overline{H}(P) \ge 0$ by Lemma 4.15. We construct a subsolution $(\varphi, \psi, 0)$ of (E_P) for $|P|$ small enough. Let

$$W_1 = (\frac{6}{16}, \frac{10}{16}),\ W_2 = (\frac{5}{16}, \frac{11}{16}),\ W_3 = (\frac{4}{16}, \frac{12}{16}).$$

Let $P < 0$ for simplicity. We define the functions φ, ψ by

$$\varphi(\xi) := \begin{cases} -P \cdot \xi & \text{for } x \in W_2 \\ 0 & \text{for } \xi \in \mathbb{T} \setminus W_3 \end{cases}$$

and $|D\varphi| \le C_1|P|$ for some $C_1 > 0$, $0 \le \varphi \le -P \cdot \xi$ on $[0, 1]$ and

$$\psi(\xi) = \begin{cases} -P \cdot \xi & \text{for } \xi \in W_1 \\ C_2 & \xi \in \mathbb{T} \setminus W_2 \end{cases}$$

for some $C_2 \in (1/64, 1)$, $|P + D\psi| \le 1$, and $\psi \ge -P \cdot \xi$ on $[0, 1]$.

We have

$$\begin{aligned}&|P+D\varphi(\xi)|^2-V_1(\xi)+\varphi(\xi)-\psi(\xi)\\ &\le\begin{cases}\varphi(\xi)-\psi(\xi)\le 0 & \text{if } \xi\in W_2\\ 2(C_1^2+1)|P|^2+\varepsilon_0+|P|-C_2 & \text{if } \xi\in\mathbb{T}\setminus W_2.\end{cases}\end{aligned}$$

If $|P|$ and ε_0 are small enough, then $|P+D\varphi(\xi)|^2-V_1(\xi)+\varphi(\xi)-\psi(\xi)\le 0$ on $\mathbb{T}$. Besides,

$$\begin{aligned}&|P+D\psi(\xi)|^2-V_2(\xi)+\psi(\xi)-\varphi(\xi)\\ &\le\begin{cases}0 & \text{if } \xi\in W_1\\ 1-2+C_2-0\le 0 & \text{if } \xi\in\mathbb{T}\setminus W_1.\end{cases}\end{aligned}$$




Fig. 4.3. The graph of φ.




Fig. 4.4. The graph of ψ.

Thus $(\varphi,\psi,0)$ is a subsolution of (E_P), and the proof is complete. □

Remark 4.16. It is worth to notice that

$$\overline{H}(0)\neq-\frac{1}{2}\min_{x\in\mathbb{T}^n}(V_1+V_2)(x)$$

in general. Indeed, set

$$\begin{aligned}V_1(\xi)&=4\pi^2\sin^2(2\pi\xi)+\cos(2\pi\xi)-\sin(2\pi\xi),\\ V_2(\xi)&=4\pi^2\cos^2(2\pi\xi)+\sin(2\pi\xi)-\cos(2\pi\xi).\end{aligned}$$

If we check that $(\cos(2\pi\xi),\sin(2\pi\xi),0)$ is a solution of (E_0), then we realize $\overline{H}(0)=0$. In this case

$$\overline{H}(0)=0\neq-2\pi^2=-\frac{1}{2}(V_1+V_2)(\xi)\text{ for all }\xi\in\mathbb{T}.$$

In Theorem 4.11 the fact that $\Pi(\mathbb{R}^n\setminus(U_1\cup U_2))$ is connected, where $U_i=\{V_i=0\}$ plays an important role in the construction of subsolutions as stated just before Theorem 4.13. In the next couple of Theorems we make new observations that we can get the flat parts of effective Hamiltonians even though $\Pi(\mathbb{R}^n\setminus(U_1\cup U_2))$ is not connected.

Theorem 4.17. *Let $n = 1$ and assume $V_1 \equiv 0$, $V_2 \geq 0$ on $[0,1]$ and $\{V_2 = 0\} = \{1/2\}$. Then there exists $\gamma > 0$ such that $\overline{H}(P) = 0$ for $|P| \leq \gamma$.*

Sketch of Proof. The proof is almost the same as the proof of Theorem 4.14 but let us present it here for the sake of clarity. Since $\min_{\xi \in \mathbb{T}^n}(V_1 + V_2)(\xi) = 0$, we have $\overline{H}(P) \geq 0$ by Lemma 4.15. Let

$$W_1 = (\frac{3}{8}, \frac{5}{8}),\ W_2 = (\frac{2}{8}, \frac{6}{8}),\ W_3 = (\frac{1}{8}, \frac{7}{8}).$$

There exists $M \in (0,1)$ so that

$$V_2(\xi) \geq M \text{ for } \xi \notin W_1.$$

Assume $P < 0$ for simplicity. We now construct the functions φ, ψ so that $(\varphi, \psi, 0)$ is a subsolution of (E_P) for small $|P|$, which implies the conclusion. Take $|P| \leq M/4$ first. We define the functions φ, ψ by

$$\varphi(\xi) := \begin{cases} -P \cdot \xi & \text{for } x \in W_2 \\ 0 & \text{for } \xi \in \mathbb{T} \setminus W_3 \end{cases}$$

and $|D\varphi| \leq C_1|P|$ for some $C_1 > 0$, $0 \leq \varphi \leq -P \cdot \xi$ on $[0,1]$ and

$$\psi(\xi) = \begin{cases} -P \cdot \xi & \text{for } \xi \in W_1 \\ C_2 & \xi \in \mathbb{T} \setminus W_2 \end{cases}$$

for some $C_2 \in (M/128, M/2)$, $|P + D\psi| \leq M/2$, and $\psi \geq -P \cdot \xi$ on $[0,1]$.

We have

$$|P + D\varphi(\xi)|^2 + \varphi(\xi) - \psi(\xi) \leq \begin{cases} \varphi(\xi) - \psi(\xi) \leq 0 & \text{if } \xi \in W_2 \\ 2(C_1^2 + 1)|P|^2 + |P| - C_2 & \text{if } \xi \in \mathbb{T} \setminus W_2. \end{cases}$$

If $|P|$ is small enough, then $|P + D\varphi(\xi)|^2 + \varphi(\xi) - \psi(\xi) \leq 0$ on $\mathbb{T}$. Besides,

$$\begin{aligned} &|P + D\psi(\xi)|^2 - V_2(\xi) + \psi(\xi) - \varphi(\xi) \\ &\leq \begin{cases} 0 & \text{if } \xi \in W_1 \\ \dfrac{M^2}{4} - M + C_2 - 0 \leq \dfrac{M^2}{4} - M + \dfrac{M}{2} \leq 0 & \text{if } \xi \in \mathbb{T} \setminus W_1. \end{cases} \end{aligned}$$

Thus $(\varphi, \psi, 0)$ is a subsolution of (E_P), and the proof is complete. □

We can actually generalize Theorem 4.17 as following.

Theorem 4.18. *Assume that $V_1 \equiv 0$, $V_2 \geq 0$ and there exist an open set W in $\mathbb{T}^n$ and a vector $q \in \mathbb{R}^n$ such that $\Pi(q + W) \Subset (0,1)^n$ and $\emptyset \neq \{V_2 = 0\} \subset W$. Then there exists $\gamma > 0$ such that $\overline{H}(P) = 0$ for $|P| \leq \gamma$.*

The proof of this Theorem is basically the same as the proof of Theorem 4.17, hence omitted. The following Corollary is a direct consequence of Theorem 4.18

Corollary 4.19. *Assume that V_1, $V_2 \geq 0$ and there exist an open set W in $\mathbb{T}^n$ and a vector $q \in \mathbb{R}^n$ such that $\Pi(q+W) \Subset (0,1)^n$ and*

$$\emptyset \neq \{V_1 = 0\} \cap \{V_2 = 0\} \subset \{V_2 = 0\} \subset W.$$

Then there exists $\gamma > 0$ such that $\overline{H}(P) = 0$ for $|P| \leq \gamma$.

The result of Corollary 4.19 is pretty surprising in the sense that flat part around 0 of $\overline{H}$ occurs even though we do not know much information about V_1. More precisely, we only need to control well $\{V_2 = 0\}$ and do not need to care about $\{V_1 = 0\}$ except that $\{V_1 = 0\} \cap \{V_2 = 0\} \neq \emptyset$.

Finally, we consider a situation in which the requirements of Theorem 4.18 and Corollary 4.19 fail.

Theorem 4.20. *We take two potentials $V^i : \mathbb{T} \to [0, \infty)$ such that V^i are continuous and $\{V^i = 0\} = \{y_{0i}\}$ for some $y_{0i} \in \mathbb{T}$ for $i = 1, 2$. Assume that $V_1(\xi_1, \xi_2) = V^1(\xi_1)$ and $V_2(\xi_1, \xi_2) = V^2(\xi_2)$ for $(\xi_1, \xi_2) \in \mathbb{T}^2$. Then there exists $\gamma > 0$ such that $\overline{H}(P) = 0$ for $|P| \leq \gamma$.*




Fig. 4.5. The figures of $\{V_i = 0\}$

Proof. By using Theorem 4.17, for $P = (P_1, P_2)$ with $|P|$ small enough, there exist two pairs $(\varphi_i, \psi_i) \in C^{0,1}(\mathbb{T})^2$ for $i = 1, 2$ such that

$$\begin{cases} |P_1 + \varphi_1'(\xi_1)|^2 - V^1(\xi_1) + \varphi_1(\xi_1) - \psi_1(\xi_1) = 0, \\ |P_1 + \psi_1'(\xi_1)|^2 + \psi_1(\xi_1) - \varphi_1(\xi_1) = 0 \end{cases}$$

and

$$\begin{cases} |P_2 + \varphi_2'(\xi_2)|^2 + \varphi_2(\xi_2) - \psi_2(\xi_2) = 0, \\ |P_2 + \psi_2'(\xi_2)|^2 - V^2(\xi_2) + \psi_2(\xi_2) - \varphi_2(\xi_2) = 0 \end{cases}$$

Now let $v_1(\xi_1, \xi_2) = \varphi_1(\xi_1) + \varphi_2(\xi_2)$, $v_2(\xi_1, \xi_2) = \psi_1(\xi_1) + \psi_2(\xi_2)$ for $(\xi_1, \xi_2) \in \mathbb{T}^2$. For $P = (P_1, P_2)$ with $|P| \leq \gamma$, we easily get that $(v_1, v_2, 0)$ is a solution of (E_P), which means $\overline{H}(P) = 0$. □

5. Generalization

In this section we consider weakly coupled systems of m-equations for $m \geq 2$

$$(u_i^\varepsilon)_t + H_i(\frac{x}{\varepsilon}, Du_i^\varepsilon) + \frac{1}{\varepsilon}\sum_{j=1}^m c_{ij}(u_i^\varepsilon - u_j^\varepsilon) = 0 \text{ in } \mathbb{R}^n \times (0,T) \text{ for } i = 1, \dots, m,$$

with

$$u_i^\varepsilon(x,0) = f_i(x) \text{ on } \mathbb{R}^n \text{ for } i = 1, \dots, m,$$

where c_{ij} are given nonnegative constants which are assumed to satisfy

$$\sum_{j=1}^m c_{ij} = 1 \text{ for all } i = 1, \dots, m. \tag{5.1}$$

Set

$$K := \begin{pmatrix} c_{11} & \cdots & c_{1m} \\ \vdots & \ddots & \vdots \\ c_{m1} & \cdots & c_{mm} \end{pmatrix}, \ \mathbf{u}^\varepsilon := \begin{pmatrix} u_1^\varepsilon \\ \vdots \\ u_m^\varepsilon \end{pmatrix}, \text{ and } \mathbf{f} := \begin{pmatrix} f_1 \\ \vdots \\ f_m \end{pmatrix}.$$

Then the problem can be written as

$$\begin{cases} \mathbf{u}_t^\varepsilon + \begin{pmatrix} H_1(x/\varepsilon, Du_1^\varepsilon) \\ \vdots \\ H_m(x/\varepsilon, Du_m^\varepsilon) \end{pmatrix} + \dfrac{1}{\varepsilon}(I-K)\mathbf{u}^\varepsilon = 0 & \text{in } \mathbb{R}^n \times (0,T), \\ \mathbf{u}^\varepsilon(\cdot,0) = \mathbf{f} & \text{on } \mathbb{R}^n, \end{cases} \tag{5.2}$$

where I is the identity matrix of size m. We obtain the following result.

Theorem 5.1. *The functions u_i^ε converge locally uniformly to the same limit u in $\mathbb{R}^n \times (0,T)$ as $\varepsilon \to 0$ for $i = 1, \dots, m$ and u solves*

$$\begin{cases} u_t + \overline{H}(Du) = 0 & \text{in } \mathbb{R}^n \times (0,T) \\ u(x,0) = \overline{f}(x) & \text{on } \mathbb{R}^n, \end{cases}$$

where $\overline{H}$ is the associated effective Hamiltonian and

$$\overline{f}(x) := \frac{1}{m}\sum_{i=1}^m f_i(x).$$

We only present barrier functions which are generalizations of the barrier function in case $m = 2$ defined by (2.1) in Lemma 2.1. Set

$$\mathbf{w}^\pm(\mathbf{x}, \mathbf{t}) := (\overline{f} \pm Ct)\mathbf{j} + \mathbf{g}^\varepsilon(x,t),$$

where C is a positive constant which will be fixed later, $\mathbf{j} := (1, \dots, 1)^T$ and

$$\mathbf{g}^\varepsilon(x,t) := \left[e^{\frac{t}{\varepsilon}(K-I)}\mathbf{h}\right](x), \ \mathbf{h}(x) := \mathbf{f}(x) - \overline{f}(x)\mathbf{j}.$$

Since we assume (5.1), we can easily check that the Frobenius root of K, i.e., the maximum of the eigenvalues of K, is 1 and moreover $\mathbf{j}$ is an associated eigenvector. Moreover by the Perron–Frobenius theorem we have

Lemma 5.2. *There exists $\delta > 0$ such that $|e^{t(K-I)}\mathbf{h}| \le e^{-\delta t}|\mathbf{h}|$ provided that $\mathbf{h} \cdot \mathbf{j} = 0$.*

See [24, Lemma 5.2] for a more general result.

Proposition 5.3. *The functions $\mathbf{w}^{\pm}$ are a subsolution and a supersolution of (5.2) with $\mathbf{w}^{\pm}(\cdot, 0) = \mathbf{f}$ on $\mathbb{R}^n$, respectively, if $C > 0$ is large enough.*

Proof. It is easy to check $\mathbf{w}^{\pm}(\cdot, 0) = \mathbf{f}$ on $\mathbb{R}^n$. Note that

$$\frac{\partial \mathbf{g}^{\varepsilon}}{\partial t} = \frac{1}{\varepsilon}(K - I)\mathbf{g}^{\varepsilon} \text{ and } |D\mathbf{g}| \le Ce^{-\frac{\delta t}{\varepsilon}}.$$

Thus, we can check easily that $\mathbf{w}^{\pm}$ are a subsolution and a supersolution of (5.2), respectively, if $C > 0$ is large enough. □

By a rather standard argument by using the perturbed test functions we can get Theorem 5.1 as in the proof of Theorem 1.2.

6. Dirichlet Problems

In this section we consider the asymptotic behavior, as ε tends to 0, of the viscosity solutions $(u_1^{\varepsilon}, u_2^{\varepsilon})$ of Dirichlet boundary problems for weakly coupled systems of Hamilton–Jacobi equations

$$(\mathrm{D}_{\varepsilon}) \qquad \begin{cases} u_1^{\varepsilon} + H_1(\dfrac{x}{\varepsilon}, Du_1^{\varepsilon}) + \dfrac{1}{\varepsilon}(u_1^{\varepsilon} - u_2^{\varepsilon}) = 0 & \text{in } \Omega, \\ u_2^{\varepsilon} + H_2(\dfrac{x}{\varepsilon}, Du_2^{\varepsilon}) + \dfrac{1}{\varepsilon}(u_2^{\varepsilon} - u_1^{\varepsilon}) = 0 & \text{in } \Omega, \\ u_i^{\varepsilon}(x) = g_i(x) & \text{on } \partial\Omega, \end{cases}$$

where Ω is a bounded domain of $\mathbb{R}^n$ with the Lipschitz boundary, the Hamiltonians $H_i \in C(\mathbb{R}^n \times \mathbb{R}^n)$ are assumed to satisfy (A1)-(A2) and $g_i \in C(\partial\Omega)$ are given functions.

Concerning the Dirichlet problem, classical works required continuous solutions up to the boundary to satisfy the prescribed data on the entire boundary. This can be achieved for special classes of equations by imposing compatibility conditions on the boundary data or by assuming the existence of appropriate super and subsolutions. However, in general, we do not expect that there exists a (viscosity) solution satisfying the boundary condition in the classical sense. After Soner studied the state constraints problems in terms of PDE, the viscosity formulation of Dirichlet conditions was introduced by Barles and Perthame [3] and Ishii [22]. In this paper we deal with solutions satisfying Dirichlet boundary conditions in the sense of viscosity solutions.

Theorem 6.1. *Let $(u_1^\varepsilon, u_2^\varepsilon)$ be the solution of (D_ε). Then u_i^ε converge locally uniformly to the same limit u on Ω as $\varepsilon \to 0$ for $i = 1, 2$ and u solves*

$$\begin{cases} u + \overline{H}(Du) = 0 & \text{in } \Omega, \\ u(x,t) = \overline{g} & \text{on } \partial\Omega, \end{cases} \tag{6.1}$$

where $\overline{g} := \min\{g_1, g_2\}$ on $\partial\Omega$.

Lemma 6.2. *If $(u_1^\varepsilon, u_2^\varepsilon)$ is a bounded uppersemicontinuous subsolution of (D_ε), then $u_i^\varepsilon(x) \le g_i(x)$ for all $x \in \partial\Omega$ and $i = 1, 2$.*

Proof. Fix $x_0 \in \partial\Omega$. Choose a sequence $\{x_k\}_{k\in\mathbb{N}} \subset \mathbb{R}^n \setminus \overline{\Omega}$ such that $|x_0 - x_k| = 1/k^2$. Define the functions $\phi_1 : \overline{\Omega} \to \mathbb{R}$ by $\phi_1(x) := u_1^\varepsilon(x) - k|x - x_k|$. Let $r > 0$ and $\xi_k \in B(x_0, r) \cap \overline{\Omega}$ be a maximum point of ϕ_1 on $B(x_0, r) \cap \overline{\Omega}$. Since $\phi_1(\xi_k) \ge \phi_1(x_0)$, we have $k|\xi_k - x_k| \le u_1^\varepsilon(\xi_k) - u_1^\varepsilon(x_0) + k|x_0 - x_k| \le C$, where $C > 0$ is a constant independent of k. Thus, $\xi_k \to x_0$ as $k \to \infty$. Moreover, noting that $u_1^\varepsilon(x_0) \le \liminf_{k\to\infty}(u_1^\varepsilon(\xi_k) + k|x_0 - x_k|) \le \limsup_{k\to\infty} u_1^\varepsilon(\xi_k) + \limsup_{k\to\infty} k|x_0 - x_k| \le u_1^\varepsilon(x_0)$, we get $u_1^\varepsilon(\xi_k) \to u_1^\varepsilon(x_0)$ as $k \to \infty$. By the viscosity property of u_1^ε, we have

$$\begin{aligned} &u_1^\varepsilon(\xi_k) + H_1\Big(\frac{\xi_k}{\varepsilon}, p_k\Big) + \frac{1}{\varepsilon}(u_1^\varepsilon(\xi_k) - u_2(\xi_k)) \le 0 \text{ or} \\ &u_1^\varepsilon(\xi_k) \le g_1(\xi_k), \end{aligned} \tag{6.2}$$

where $p_k = k(\xi_k - x_k)/|\xi_k - x_k|$. Noting that $|p_k| = k$, by (A1), we see that the left-hand side of (6.2) is positive for a sufficiently large $k \in \mathbb{N}$ and then we must have $u_1^\varepsilon(\xi_k) \le g_1(\xi_k)$. Sending $k \to \infty$, we get $u_1^\varepsilon(x_0) \le g_1(x_0)$. Similaly, we get $u_2^\varepsilon(x_0) \le g_2(x_0)$ on $\partial\Omega$. □

Lemma 6.3. *The families $\{u_i^\varepsilon\}_{\varepsilon>0}$ are equi-Lipschitz continuous in Ω for $i = 1, 2$.*

Proof. Set $M := \max_{i=1,2}(\|H_i(\cdot, 0)\|_{L^\infty(\mathbb{R}^n)} + \|g_i\|_{L^\infty(\partial\Omega)})$. Then $(-M, -M)$ and (M, M) are a subsolution and a supersolution of (D_ε), respectively. By the comparison principle for (D_ε) we have $|u_i^\varepsilon| \le M$. Adding two equations in (D_ε) we get

$$u_1^\varepsilon + u_2^\varepsilon + H_1\Big(\frac{x}{\varepsilon}, Du_1^\varepsilon\Big) + H_2\Big(\frac{x}{\varepsilon}, Du_2^\varepsilon\Big) = 0$$

for almost every $x \in \Omega$, which implies that $|Du_i^\varepsilon| \le M'$ in the sense of viscosity solutions for some $M' > 0$, which is independent of ε. □

Proof of Theorem 1.2. By Lemma 6.3 we can extract a subsequence $\{\varepsilon_j\}$ converging to 0 so that $u_i^{\varepsilon_j}$ converges locally uniformly to $u_i \in C(\overline{\Omega})$ for $i = 1, 2$. By usual observations, we get that $u_1 = u_2 =: u$. Since (6.1) has a unique solution, it is enough for us to prove that u is a solution of (6.1).

We only prove that u is a supersolution of (6.1), since in view of Lemma 6.2 we can easily see that u is a subsolution of (6.1).

Let $\phi \in C^1(\overline{\Omega})$ be a test function such that $u - \phi$ takes a strict minimum at $x_0 \in \overline{\Omega}$. We only consider the case where $x_0 \in \partial\Omega$, since we can prove by a similar way to the proof of Theorem 1.2 in the case where $x_0 \in \Omega$. It is enough for us to prove that $u(x_0) + \overline{H}(D\phi(x_0)) \geq 0$ provided that $(u - \overline{g})(x_0) < 0$.

Let (v_1, v_2) be a solution of (E_P) with $P := D\phi(x_0)$. We consider

$$m^\varepsilon := \min_{i\in\{1,2\}} \min_{x\in\overline{\Omega}} \big(u_i^\varepsilon(x) - \phi(x) - \varepsilon v_i(\frac{x}{\varepsilon})\big).$$

Pick $i^\varepsilon \in \{1,2\}$ and $x^\varepsilon \in \overline{\Omega}$ so that $m^\varepsilon = u_{i^\varepsilon}^\varepsilon(x^\varepsilon) - \phi(x^\varepsilon) - \varepsilon v_{i^\varepsilon}(x^\varepsilon/\varepsilon)$. Also choose $j^\varepsilon \in \{1,2\}$ such that $\{i^\varepsilon, j^\varepsilon\} = \{1,2\}$. We only consider the case where $x^\varepsilon \in \partial\Omega$ again. Since u_{i_ε} converges to u locally uniformly on $\overline{\Omega}$, $\varepsilon v(\cdot/\varepsilon)$ converges to 0 uniformly on $\overline{\Omega}$ as $\varepsilon \to 0$ and $u - \phi$ takes a strict maximum at x_0, we see that $x^\varepsilon \to x_0$ as $\varepsilon \to 0$. Thus, if ε is small enough, then we may assume that $(u_{i^\varepsilon}^\varepsilon - g_{i^\varepsilon})(x^\varepsilon) < 0$.

For $\alpha > 0$ we define the function $\Phi_\alpha : \overline{\Omega} \times \mathbb{R}^n \to \mathbb{R}$ by

$$\Phi_\alpha(x,y) := u_{i^\varepsilon}^\varepsilon(x) - \phi(x) - \varepsilon v_{i^\varepsilon}\big(\frac{y}{\varepsilon}\big) + \frac{1}{2\alpha^2}|x-y|^2 + \frac{1}{2}|x - x^\varepsilon|^2.$$

Let Φ_α achieve its minimum over $\overline{\Omega}\times\mathbb{R}^n$ at some $(x_\alpha^\varepsilon, y_\alpha^\varepsilon)$. Since we may assume by taking a subsequence if necessary that $x_\alpha^\varepsilon \to x^\varepsilon$ as $\alpha \to 0$, we have

$$(u_{i^\varepsilon} - g_{i^\varepsilon})(x_\alpha^\varepsilon) < 0 \text{ for small } \alpha > 0.$$

Therefore, by the definition of viscosity solutions, we have

$$u_{i^\varepsilon} + H_{i^\varepsilon}(\frac{x_\alpha^\varepsilon}{\varepsilon}, D\phi(x_\alpha^\varepsilon) - p_\alpha^\varepsilon - (x_\alpha^\varepsilon - x^\varepsilon)) + \frac{1}{\varepsilon}(u_{i^\varepsilon}^\varepsilon - u_{j^\varepsilon}^\varepsilon)(x_\alpha^\varepsilon) \geq 0,$$

where $p_\alpha^\varepsilon := (x_\alpha^\varepsilon - y_\alpha^\varepsilon)/\alpha^2$. Also, we have

$$H_i(\frac{y_\alpha^\varepsilon}{\varepsilon}, P - p_\alpha^\varepsilon) + (v_{i^\varepsilon} - v_{j^\varepsilon})(\frac{y_\alpha^\varepsilon}{\varepsilon}) \leq \overline{H}(P),$$

since (v_1, v_2) is a solution of (E_P).

A priori Lipschitz estimate implies $|p_\alpha^\varepsilon| \leq C$ for some $C > 0$ which is independent of α and ε. Without loss of generality, we may assume that $p_\alpha^\varepsilon \to p^\varepsilon$ by taking a subsequence $\{\alpha_j\}$ converging to 0 if necessary. Send $\alpha \to 0$ in the above inequalities to obtain

$$\begin{aligned} &u_{i^\varepsilon}^\varepsilon(x^\varepsilon) + H_{i^\varepsilon}(\frac{x^\varepsilon}{\varepsilon}, D\phi(x^\varepsilon) - p^\varepsilon) + \frac{1}{\varepsilon}(u_{i^\varepsilon}^\varepsilon(x^\varepsilon) - u_{j^\varepsilon}^\varepsilon(x^\varepsilon)) \geq 0, \\ &H_{i^\varepsilon}(\frac{x^\varepsilon}{\varepsilon}, P - p^\varepsilon) + v_{i^\varepsilon}(\frac{x^\varepsilon}{\varepsilon}) - v_{j^\varepsilon}(\frac{x^\varepsilon}{\varepsilon}) \leq \overline{H}(P). \end{aligned}$$

Noting that $u_{i^\varepsilon}^\varepsilon(x^\varepsilon) - \phi(x^\varepsilon) - \varepsilon v_{i^\varepsilon}(x^\varepsilon/\varepsilon) \leq u_{j^\varepsilon}^\varepsilon(x^\varepsilon) - \phi(x^\varepsilon) - \varepsilon v_{j^\varepsilon}(x^\varepsilon/\varepsilon)$, we get that

$$u_{i^\varepsilon}^\varepsilon(x^\varepsilon) + \overline{H}(P) \geq H_{i^\varepsilon}(\frac{x^\varepsilon}{\varepsilon}, D\phi(x^\varepsilon) - p^\varepsilon) - H_{i^\varepsilon}(\frac{x^\varepsilon}{\varepsilon}, P - p^\varepsilon) \geq -\sigma(|D\phi(x^\varepsilon) - P|).$$

Sending $\varepsilon \to 0$ yields the conclusion. □

In order to explain the relation between (D_ε) and the exit-time problem in the optimal control theory, we assume that the Hamiltonians H_i are convex in the p-variable henceforth. We next define the associated value functions, which give us an intuition about the effective boundary datum $\overline{g}$ in Theorem 6.1.

For $\varepsilon > 0$ we define the functions $u_i^\varepsilon : \overline{\Omega} \to \mathbb{R}$ by

$$u_i^\varepsilon(x) := \inf\Big\{\mathbb{E}_i\Big(\int_0^\tau e^{-s} L_{\nu^\varepsilon(s)}\big(\frac{\eta(s)}{\varepsilon}, -\dot{\eta}(s)\big)\, ds + e^{-\tau} g_{\nu^\varepsilon(\tau)}(\eta(\tau))\Big)\Big\}, \tag{6.3}$$

where the infimum is taken over $\eta \in \mathrm{AC}\,([0,\infty), \overline{\Omega})$ such that $\eta(0) = x$ and $\tau \in [0,\infty]$ such that $\eta(\tau) \in \partial\Omega$ and if $\tau = \infty$, then we set $e^{-\infty} := 0$. Here $\mathbb{E}_i$ denotes the expectation of a process with $\nu^\varepsilon(0) = i$, where ν^ε is a $\{1,2\}$-valued continuous-time Markov chain given by (1.2).

Theorem 6.4. *Assume that the functions u_i^ε given by* (6.3) *are continuous on $\overline{\Omega}$. Then the pair $(u_1^\varepsilon, u_2^\varepsilon)$ is a solution of* (D_ε).

The proof of Theorem 6.4 is given in Appendix. See [3, 22] for single equations. The value functions defined by (6.3) give us an intuitive explanation of the reason why the boundary datum $\overline{g}$ of the limit solution u is the minimum of g_i for $i = 1, 2$. If we send ε to 0, then the switching rate becomes very fast but it does not really affect the exit time as we can choose to stay in $\overline{\Omega}$ as long as we like. And hence, we can control the exit state in such a way that the exit cost is the minimum of two given exit costs g_i. On the other hand, when we consider the value function (1.1) associated with the initial value problem, we cannot control the terminal state and also the timing of jumps, which are only determined by a probabilistic way given by (1.2). This is the main difference between Dirichlet problems and initial value problems and the reason why the effective Dirichlet boundary value and the effective initial value are different.

7. Appendix

We now prove Theorems 7.1, and 6.4 by basically using the dynamic programming principles, which are pretty standard in the theory of viscosity solutions. Throughout this section we always assume in addition to (A1), (A2) that $p \mapsto H_i(\xi, p)$ are convex for $i = 1, 2$.

Theorem 7.1 (Verification Theorem)**.** *Assume that the functions u_i^ε given by* (1.1) *are continuous on $\mathbb{R}^n \times [0,T]$. Then the pair $(u_1^\varepsilon, u_2^\varepsilon)$ is a solution of* (C_ε).

Let $\varepsilon = 1$ for simplicity in what follows. By abuse of notations we write (u_1, u_2) for (u_1^1, u_2^1).

Proposition 7.2 (Dynamic Programming Principle)**.** *For any $x \in \mathbb{R}^n$, $0 \le h \le t$ and $i = 1, 2$ we have*

$$u_i(x,t) = \inf\Big\{\mathbb{E}_i\Big(\int_0^h L_{\nu(s)}(\eta(s), -\dot\eta(s))\,ds + u_{\nu(h)}(\eta(h), t-h)\Big)\Big\}, \tag{7.1}$$

where the infimum is taken over $\eta \in \mathrm{AC}([0,h],\mathbb{R}^n)$ with $\eta(0) = x$.

Proof. We denote by $v_i(x,t;h)$ the right-hand side of (7.1). Let η be a trajectory in $\mathrm{AC}([0,t],\mathbb{R}^n)$ with $\eta(0) = x$ and ν be a process with $\nu(0) = i$ which satisfies (1.2). Set $\tilde\eta(s) := \eta(s+h)$ and $\tilde\nu(s) := \nu(s+h)$ for $s \in [0, t-h]$. We have

$$\begin{aligned}
&\mathbb{E}_i\Big(\int_0^t L_{\nu(s)}(\eta(s), -\dot\eta(s))\,ds + f_{\nu(t)}(\eta(t))\Big)\\
&= \mathbb{E}_i\Big(\int_0^h L_{\nu(s)}(\eta(s), -\dot\eta(s))\,ds + \int_h^t L_{\nu(s)}(\eta(s), -\dot\eta(s))\,ds + f_{\nu(t)}(\eta(t))\Big)\\
&= \mathbb{E}_i\Big(\int_0^h L_{\nu(s)}(\eta(s), -\dot\eta(s))\,ds\Big) + \mathbb{E}_i\Big(\int_0^{t-h} L_{\tilde\nu(s)}(\tilde\eta(s), -\dot{\tilde\eta}(s))\,ds + f_{\tilde\nu(t-h)}(\tilde\eta(t-h))\Big)\\
&\ge \mathbb{E}_i\Big(\int_0^h L_{\nu(s)}(\eta(s), -\dot\eta(s))\,ds + u_{\nu(h)}(\eta(h), t-h)\Big)\\
&\ge v_i(x,t;h),
\end{aligned}$$

which implies $u_i(x,t) \ge v_i(x,t;h)$.

Let $\delta_1 \in \mathrm{AC}([0,h],\mathbb{R}^n)$ and $\delta_2 \in \mathrm{AC}([0,t-h],\mathbb{R}^n)$ be trajectories with $\delta_1(h) = \delta_2(0)$ and $\delta_1(0) = x$. Set

$$\eta(s) := \begin{cases} \delta_1(s) & \text{for all } s \in [0,h],\\ \delta_2(s-h) & \text{for all } s \in [h,t]. \end{cases}$$

Let ν be a process with $\nu(0) = i$ which satisfies (1.2). Note that

$$\begin{aligned}
&\int_0^h L_{\nu(s)}(\delta_1(s), -\dot\delta_1(s))\,ds + \int_0^{t-h} L_{\nu(s+h)}(\delta_2(s), -\dot\delta_2(s))\,ds + f_{\nu(t)}(\delta_2(t-h))\\
&= \int_0^t L_{\nu(s)}(\eta(s), -\dot\eta(s))\,ds + f_{\nu(t)}(\eta(t)).
\end{aligned}$$

We have

$$\begin{aligned}
&\mathbb{E}_i\Big(\int_0^h L_{\nu(s)}(\delta_1(s), -\dot\delta_1(s))\,ds + \int_0^{t-h} L_{\nu(s+h)}(\delta_2(s), -\dot\delta_2(s))\,ds + f_{\nu(t)}(\delta_2(t-h))\Big)\\
&= \mathbb{E}_i\Big(\int_0^t L_{\nu(s)}(\eta(s), -\dot\eta(s))\,ds + f_{\nu(t)}(\eta(t))\Big)\\
&\ge u_i(x,t).
\end{aligned}$$

Take the infimum on all admissible δ_2 to obtain

$$\mathbb{E}_i\Big(\int_0^h L_{\nu(s)}(\delta_1(s), -\delta_1(s))\,ds + u_{\nu(h)}(\delta_1(h), t-h)\Big) \geq u_i(x,t),$$

which implies $v_i(x,t;h) \geq u_i(x,t)$. □

Proof of Theorem 7.1. It is obvious to see that $(u_1,u_2)(\cdot,0) = (f_1,f_2)$ on $\mathbb{R}^n$. We first prove that u_1 is a subsolution of (C_1). We choose a function $\phi \in C^1(\mathbb{R}^n \times (0,T))$ such that $u_1 - \phi$ has a strict maximum at $(x_0,t_0) \in \mathbb{R}^n \times (0,T)$ and $(u_1-\phi)(x_0,t_0) = 0$.

Let $h > 0$. By Proposition 7.1 we have

$$u_1(x_0,t_0) \leq \mathbb{E}_i\Big(\int_0^h L_{\nu(s)}(\eta(s), -\dot\eta(s))\,ds + u_{\nu(h)}(\eta(h), t_0-h)\Big) \tag{7.2}$$

for any $\eta \in \mathrm{AC}\,([0,h],\mathbb{R}^n)$ with $\eta(0) = x_0 \in \mathbb{R}^n$ and $\dot\eta(0) = q \in \mathbb{R}^n$. Since ν is a continuous-time Markov chain which satisfies (1.2), the probability that $\nu(h) = 2$ is $c_1h + o(h)$ and the probability that $\nu(h) = 1$ is $1 - (c_1h + o(h))$. By (7.2) we obtain

$$\begin{aligned}
&\phi(x_0,t_0) = u_1(x_0,t_0)\\
&\leq (1 - c_1h - o(h))\Big(\int_0^h L_1(\eta,-\dot\eta)\,ds + u_1(\eta(h),t_0-h)\Big)\\
&\quad + (c_1h + o(h))\Big(\int_0^h L_2(\eta,-\dot\eta)\,ds + u_2(\eta(h),t_0-h)\Big)\\
&\leq \int_0^h L_1(\eta,-\dot\eta)\,ds + \phi(\eta(h),t_0-h)\\
&\quad + (c_1h+o(h))\Big(\int_0^h L_2(\eta,-\dot\eta)\,ds + u_2(\eta(h),t_0-h) - \int_0^h L_1(\eta,-\dot\eta)\,ds - u_1(\eta(h),t_0-h)\Big).
\end{aligned}$$

Thus,

$$\begin{aligned}
&\frac{\phi(\eta(0),t_0) - \phi(\eta(h),t_0-h)}{h}\\
&\leq \frac{1}{h}\int_0^h L_1(\eta,-\dot\eta)\,ds + (c_1 + \frac{o(h)}{h})(u_2(\eta(h),t_0-h) - u_1(\eta(h),t_0-h))\\
&\quad + (c_1 + \frac{o(h)}{h})\Big(\int_0^h L_2(\eta,-\dot\eta)\,ds - \int_0^h L_1(\eta,-\dot\eta)\,ds\Big).
\end{aligned}$$

Sending $h \to 0$, we obtain

$$\phi_t(x_0,t_0) + D\phi(x_0,t_0)\cdot(-q) \leq L_1(x_0,-q) + c_1(u_2-u_1)(x_0,t_0) \text{ for all } q \in \mathbb{R}^n,$$

which implies $\phi_t(x_0,t_0) + H(x_0, D\phi(x_0,t_0)) + c_1(u_2-u_1)(x_0,t_0) \leq 0$.

Next we prove that u_1 is a supersolution of (C_1). We choose a function $\phi \in C^1(\mathbb{R}^n \times (0,T))$ such that $u_1 - \phi$ has a strict minimum at $(x_0,t_0) \in \mathbb{R}^n \times (0,T)$ and $(u_1-\phi)(x_0,t_0) = 0$. Take $h, \delta > 0$. By Proposition 7.1 we have

$$u_1(x_0,t_0) + \delta > \mathbb{E}_i\Big(\int_0^h L_{\nu(s)}(\eta_\delta(s), -\dot\eta_\delta(s))\, ds + u_{\nu(h)}(\eta_\delta(h), t_0 - h)\Big) \tag{7.3}$$

for some $\eta_\delta \in \mathrm{AC}\,([0,h],\mathbb{R}^n)$ with $\eta_\delta(0) = x_0$. Since ν is a continuous-time Markov chain which satisfies (1.2), by a similar calculation to the above we obtain

$$\begin{aligned}
&\phi(x_0,t_0) + \delta = u_1(x_0,t_0) + \delta \\
&> \int_0^h L_1(\eta, -\dot\eta)\, ds + \phi(\eta(h), t_0 - h) \\
&\quad + (c_1 h + o(h))\Big(\int_0^h L_2(\eta, -\dot\eta)\, ds + u_2(\eta(h), t_0 - h) - \int_0^h L_1(\eta, -\dot\eta)\, ds - u_1(\eta(h), t_0 - h)\Big).
\end{aligned}$$

Thus,

$$\begin{aligned}
\frac{\delta}{h} &> \frac{1}{h}\int_0^h \frac{d\phi(\eta_\delta(s), t_0 - s)}{ds} + L_1(\eta_\delta, -\dot\eta_\delta)\, ds \\
&\quad + (c_1 + \frac{o(h)}{h})(u_2(\eta_\delta(h), t_0 - h) - u_1(\eta_\delta(h), t_0 - h)) \\
&\quad + (c_1 + \frac{o(h)}{h})\Big(\int_0^h L_2(\eta_\delta, -\dot\eta_\delta)\, ds - \int_0^h L_1(\eta_\delta, -\dot\eta_\delta)\, ds\Big) \\
&= \frac{1}{h}\int_0^h -\phi_t(\eta_\delta(s), t_0 - s) - D\phi \cdot (-\dot\eta_\delta(s)) + L_1(\eta_\delta, -\dot\eta_\delta)\, ds \\
&\quad + (c_1 + \frac{o(h)}{h})(u_2(\eta_\delta(h), t_0 - h) - u_1(\eta_\delta(h), t_0 - h)) + O(h) \\
&\geq \frac{1}{h}\int_0^h -\big(\phi_t(\eta_\delta(s), t_0 - s) + H_1(\eta_\delta(s), D\phi)\big)\, ds \\
&\quad + (c_1 + \frac{o(h)}{h})(u_2(\eta_\delta(h), t_0 - h) - u_1(\eta_\delta(h), t_0 - h)) + O(h).
\end{aligned}$$

We finally set $\delta = h^2$ and let $h \to 0$ to yield the conclusion. □

By a similar argument to the proof of Proposition 7.2 we can prove

Proposition 7.3 (Dynamic Programming Principle for (6.3))**.** *For any $x \in \mathbb{R}^n$, $h \geq 0$ and $i = 1, 2$ we have*

$$\begin{aligned}
u_i(x) = \inf\Big\{\mathbb{E}_i\Big(\int_0^{h\wedge\tau} e^{-s} L_{\nu(s)}(\frac{\eta(s)}{\varepsilon}, -\dot\eta(s))\, ds \\
+ \mathbf{1}_{\{h<\tau\}} e^{-h} u_{\nu(h)}(\eta(h)) + \mathbf{1}_{\{h\geq\tau\}} e^{-\tau} g_{\nu(\tau)}(\eta(\tau))\Big)\Big\},
\end{aligned} \tag{7.4}$$

where ν with $\nu(0) = i$ is a $\{1,2\}$-valued continuous-time Markov chain which satisfies (1.2) *and the infimum is taken over $\eta \in \mathrm{AC}([0,h],\overline{\Omega})$ such that $\eta(0) = x$ and $\tau \in [0,h]$ such that $\eta(\tau) \in \partial\Omega$.*

Proof of Theorem 6.4. We only prove in what follows that u_i satisfy the Dirichlet boundary condition in the sense of viscosity solutions, as we can prove u_i satisfy the equations by an argument similar to the proof of Theorem 7.1. Since it is clear to see that $u_i \le g_i$ on $\partial\Omega$ in the classical sense from the definition of u_i, we only need to prove that (u_1, u_2) is a supersolution of (D_ε) and particularly that u_1 satisfies the boundary condition in the viscosity solution sense. Take $x_0 \in \partial\Omega$ and suppose that

$$(u_1 - g_1)(x_0) < 0. \tag{7.5}$$

Let $\phi \in C^1(\overline{\Omega})$ satisfy $(u_1 - \phi)(x_0) = \max_{\overline{\Omega}}(u_1 - \phi) = 0$. By Proposition 7.3 we have

$$\begin{aligned}
&u_1(x_0) + h^2 \\
&> \mathbb{E}_i\Big(\int_0^{h\wedge\tau_h} e^{-s} L_{\frac{\nu(s)}{\varepsilon}}(\eta_h(s), -\dot\eta_h(s))\, ds + \mathbf{1}_{\{h<\tau_h\}} e^{-h} u_{\nu(h)}(\eta_h(h)) + \mathbf{1}_{\{h\ge\tau_h\}} e^{-\tau_h} g_{\nu(\tau_h)}(\eta_h(\tau_h))\Big) \\
&\ge \mathbb{E}_i\Big(\int_0^{h\wedge\tau_h} e^{-s} L_{\nu(s)}(\frac{\eta_h(s)}{\varepsilon}, -\dot\eta_h(s))\, ds + e^{-(h\wedge\tau_h)} u_{\nu(h\wedge\tau_h)}(\eta_h(h\wedge\tau_h))\Big)
\end{aligned}$$

for some $\eta_h \in \mathrm{AC}([0,h],\overline{\Omega})$ such that $\eta(0) = x$ and $\tau_h \in [0,h]$. In view of (7.5), we have $\tau_h > 0$ for small $h > 0$. Therefore by a similar calculation as in the proof of Theorem 7.1 we get

$$u_1(x_0) + H_1(\frac{x_0}{\varepsilon}, D\phi(x_0)) + \frac{1}{\varepsilon}(u_1 - u_2)(x_0) \ge 0. \qquad \square$$

References

1. S. N. Armstrong, P. E. Souganidis, *Stochastic homogenization of Hamilton–Jacobi and degenerate Bellman equations in unbounded environments*, J. Math. Pures Appl., to appear.
2. S. N. Armstrong, P. E. Souganidis, *Concentration phenomena for neutronic multigroup diffusion in random environments*, submitted.
3. G. Barles, B. Perthame, *Exit time problems in optimal control and vanishing viscosity method*, SIAM J. Control Optim. **26** (1988), no. 5, 1133–1148.
4. F. Cagnetti, D. Gomes, H. V. Tran, *Adjoint methods for obstacle problems and weakly coupled systems of PDE*, submitted.
5. F. Camilli, A. Cesaroni, C. Marchi, *Homogenization and vanishing viscosity in fully nonlinear elliptic equations: rate of convergence estimates*, Adv. Nonlinear Stud., **11** (2011), no. 2, 405–428.
6. F. Camilli, O. Ley, P. Loreti, *Homogenization of monotone systems of Hamilton–Jacobi equations*, ESAIM Control Optim. Calc. Var. **16** (2010), no. 1, 58–76.
7. F. Camilli, O. Ley, P. Loreti, V. Nguyen, *Large time behavior of weakly coupled systems of first-order Hamilton-Jacobi equations*, NoDEA Nonlinear Differential Equations Appl. to appear.
8. F. Camilli, C. Marchi, *Continuous dependence estimates and homogenization of quasi-monotone systems of fully nonlinear second order parabolic equations*, preprint.

9. I. Capuzzo-Dolcetta, H. Ishii, *On the rate of convergence in homogenization of Hamilton–Jacobi equations*, Indiana Univ. Math. J. **50** (2001), no. 3, 1113–1129.
10. G. Contreras, R. Iturriaga, G. P. Paternain, M. Paternain, *Lagrangian graphs, minimizing measures and Mañé's critical values*, Geom. Funct. Anal. **8** (1998), no. 5, 788–809.
11. M. C. Concordel, *Periodic homogenization of Hamilton–Jacobi equations: additive eigenvalues and variational formula*, Indiana Univ. Math. J. **45** (1996), no. 4, 1095–1117.
12. M. C. Concordel, *Periodic homogenisation of Hamilton–Jacobi equations. II. Eikonal equations*, Proc. Roy. Soc. Edinburgh Sect. A **127** (1997), no. 4, 665–689.
13. M. H. A. Davis, *Piecewise-deterministic Markov processes: a general class of nondiffusion stochastic models*, J. Roy. Statist. Soc. Ser. B **46** (1984), no. 3, 353–388.
14. A. Eizenberg, M. Freidlin, *On the Dirichlet problem for a class of second order PDE systems with small parameter*, Stochastics Stochastics Rep., **33** (3-4):111–148, 1990.
15. H. Engler, S. M. Lenhart, *Viscosity solutions for weakly coupled systems of Hamilton–Jacobi equations*, Proc. London Math. Soc. (3) **63** (1991), no. 1, 212–240.
16. L. C. Evans, *The perturbed test function method for viscosity solutions of nonlinear PDE*, Proc. Roy. Soc. Edinburgh Sect. A **111** (1989), no. 3-4, 359–375.
17. L. C. Evans, *Periodic homogenisation of certain fully nonlinear partial differential equations*, Proc. Roy. Soc. Edinburgh Sect. A **120** (1992), no. 3-4, 245–265.
18. L. C. Evans, D. Gomes, *Effective Hamiltonians and averaging for Hamiltonian dynamics. I*, Arch. Ration. Mech. Anal. **157** (2001), no. 1, 1–33.
19. B. Fehrman, *Stochastic homogenization of monotone systems of viscous Hamilton–Jacobi equations with convex nonlinearities*, preprint.
20. D. Gomes, *A stochastic analogue of Aubry-Mather theory*, Nonlinearity **15**, (2002), no. 3, 581–603.
21. K. Horie, H. Ishii, *Homogenization of Hamilton–Jacobi equations on domains with small scale periodic structure*, Indiana Univ. Math. J. **47** (1998), no. 3, 1011–1058.
22. H. Ishii, *A boundary value problem of the Dirichlet type for Hamilton–Jacobi equations*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **16** (1989), no. 1, 105–135.
23. H. Ishii, S. Koike, *Viscosity solutions for monotone systems of second-order elliptic PDEs*, Comm. Partial Differential Equations **16** (1991), no. 6-7, 1095–1128.
24. H. Ishii, K. Shimano, *Asymptotic analysis for a class of infinite systems of first-order PDE: nonlinear parabolic PDE in the singular limit*, Comm. Partial Differential Equations **28** (2003), no. 1-2, 409–438.
25. E. Kosygina, F. Rezakhanlou, S. R. S. Varadhan, *Stochastic homogenization of Hamilton–Jacobi–Bellman equations*, Comm. Pure Appl. Math. **59** (10) (2006) 1489–1521.
26. S. M. Lenhart, N. Yamada, *Viscosity solutions associated with switching game for piecewise- deterministic processes*, Stochastics Stochastics Rep., **38** (1): 27–47, 1992.
27. P.-L. Lions, G. Papanicolaou, S. R. S. Varadhan, *Homogenization of Hamilton–Jacobi equations*, unpublished work (1987).
28. P.-L. Lions, P. E. Souganidis, *Correctors for the homogenization of Hamilton–Jacobi equations in the stationary ergodic setting*, Comm. Pure Appl. Math. **56** (10) (2003) 1501–1524.
29. P.-L. Lions, P. E. Souganidis, *Stochastic homogenization of Hamilton–Jacobi and "viscous" Hamilton–Jacobi equations with convex nonlinearities–revisited*, Commun. Math. Sci. **8** (2) (2010) 627–637.
30. H. Mitake, H. V. Tran, *Remarks on the large time behavior of viscosity solutions of quasi-monotone weakly coupled systems of Hamilton–Jacobi equations*, Asymptot. Anal., to appear.
31. F. Rezakhanlou, J. E. Tarver, *Homogenization for stochastic Hamilton–Jacobi equations*, Arch. Ration. Mech. Anal. **151** (4) (2000) 277–309.

32. R. W. Schwab, *Stochastic homogenization of Hamilton–Jacobi equations in stationary ergodic spatio-temporal media*, Indiana Univ. Math. J. **58** (2) (2009) 537–581.
33. K. Shimano, *Homogenization and penalization of functional first-order PDE*, NoDEA Nonlinear Differential Equations Appl. **13** (2006), no. 1, 1–21.
34. P. E. Souganidis, *Stochastic homogenization of Hamilton–Jacobi equations and some applications*, Asymptot. Anal. **20** (1) (1999) 1–11.
35. H. V. Tran, *Adjoint methods for static Hamilton-Jacobi equations*, Calc. Var. Partial Differential Equations **41** (2011), no. 3-4, 301–319.

(H. Mitake) Department of Applied Mathematics, Faculty of Science, Fukuoka University, Fukuoka 814-0180, Japan
E-mail address: mitake@math.sci.fukuoka-u.ac.jp

(H. V. Tran) Department of Mathematics, University of California, Berkeley, CA 94720, USA
E-mail address: tvhung@math.berkeley.edu